# Strain Rates along the Alpine-Himalayan Belt from a Comprehensive GNSS Velocity Field

N. Castro-Perdomo<sup>1</sup>, S. Jónsson<sup>2</sup>, Y. Klinger<sup>3</sup>, F. Masson<sup>4</sup>,

T. W. Becker<sup>5,6,7</sup>, K. Johnson<sup>1</sup>

Indiana University, Department of Earth and Atmospheric Sciences, Bloomington, Indiana 47405, U.S.A

Physical Science and Engineering Division, King Abdullah University of Science and Technology

(KAUST), Thuwal, 23955, Saudi Arabia

Université Paris Cité, Institut de Physique du Globe de Paris, CNRS, F-75005, Paris, France

Institute for Geophysics, Jackson School of Geosciences, The University of Texas at Austin, 78758,

Austin, U.S.A

Department of Earth and Planetary Sciences, Jackson School of Geosciences, The University of Texas at Austin, 78758, Austin, U.S.A

Austin, 78758, Austin, U.S.A

Austin, U.S.A

Austin, U.S.A

Austin, U.S.A

#### **Key Points:**

16

17

19

21

22

- We build a comprehensive GNSS velocity field for the Alpine-Himalayan belt, including over 11,000 horizontal velocities
- We estimate strain rates and posterior uncertainties in the eastern Mediterranean and the India-Asia collision zone
- We find consistency between the orientation and style of geodetic and seismic strain rate tensors across the Alpine-Himalayan belt

Corresponding author: N. Castro-Perdomo, jcastrop@iu.edu

#### Abstract

23

24

25

26

27

28

29

30

32

33

34

36

37

40

41

42

43

44

45

47

48

49

50

51

52

53

57

58

59

60

61

62

63

64

65

69

70

71

The Alpine-Himalayan belt is one of Earth's most dynamic and complex regions, characterized by intense tectonic deformation and seismicity. Comprehensive analyses of continental-scale crustal deformation and seismic hazards along this extensive orogenic belt require the compilation of large geodetic datasets. In this study, we integrate 42 published Global Navigation Satellite System (GNSS) velocity fields, building an internally consistent dataset for the entire belt, spanning from Iberia to Southeast Asia and comprising 11,177 horizontal and 3,940 vertical velocities. We use this unified GNSS velocity field to estimate surface strain rates and their posterior uncertainties in the eastern Mediterranean region and the India-Asia collision zone. Our results show large-scale agreement between the orientation and style of geodetic and seismic strain rate tensors across the belt. Additionally, our analyses substantiate previously documented azimuthal alignments between principal strain rate directions and seismic anisotropy orientations, often used as a proxy for finite strain in the convecting mantle. These correlations are particularly apparent in the Aegean, North Anatolia, Tibet, Tian Shan, Altai, Sayan, and Baikal regions, underscoring the need for future research on the relationship between mantle flow and lithospheric deformation.

# Plain Language Summary

The Alpine-Himalayan Belt, extending from Iberia to Southeast Asia, is one of the most tectonically active regions on Earth, frequently affected by large and damaging earth-quakes. In this study, we compile and merge geodetic data across this vast region to construct a unified and comprehensive crustal deformation velocity field. Using this new model, we estimate the rates at which the Earth's surface deforms due to tectonic and geodynamic processes. Our results show strong agreement between the orientation and style of geodetic strain accumulation in the crust and strain release during earthquakes. Additionally, we substantiate the regional angular alignments between surface deformation patterns and azimuthal seismic anisotropy, often used as a proxy for deformation in the Earth's upper mantle.

# 1 Introduction

The Alpine-Himalayan belt, associated with the closure of the Tethyan ocean, extends over 10,000 km from Gibraltar to the Indochina Peninsula in Southeast Asia and ranks among Earth's most seismically active and tectonically complex regions (e.g., Argand, 1922; Carey, 1955; Molnar & Tapponnier, 1975; Şengör, 1986; Hatzfeld & Molnar, 2010; Reilinger & McClusky, 2011). Formed through the northward subduction of the Neo-Tethys oceanic lithosphere from the Late Mesozoic to Early Cenozoic, this extensive convergence zone has been a major collisional boundary where the Eurasian Plate converged with Gondwanaland-derived fragments, including the Nubian, Arabian, and Indian Plates, as well as several smaller lithospheric blocks (e.g., Read & Watson, 1975; Dercourt et al., 1986).

These tectonic interactions have produced a broad deformation zone along the southern margin of the Eurasian plate. This zone features a nascent convergent boundary in Northern Africa (Billi et al., 2023), active subduction along the Calabria, Hellenic, Cyprus, Makran and Sunda megathrusts (e.g., Faccenna et al., 2011; Faccenna, Becker, Auer, et al., 2014; L. Jolivet et al., 2013; Barbot & Weiss, 2021; Lindsey et al., 2023), continental collision along the Caucasus, Zagros, and Himalayas, and lateral transport zones in Anatolia and the eastern and western margins of the Tibetan Plateau (e.g., Tapponnier et al., 1982; Hatzfeld & Molnar, 2010; M. Wang & Shen, 2020). The Alpide belt also hosts prominent mountain ranges, including the Atlas, Alps, Hellenides, Taurides, Zagros, Makran, and the Himalayas, all of which experience ongoing crustal deformation and heightened

seismicity (e.g., Jackson & McKenzie, 1984; D. McKenzie, 1978; Bilham, 2006; Reilinger & McClusky, 2011), as illustrated in Fig. 1.

Renewed interest in seismic hazards along the Alpine-Himalayan belt has been spurred by a series of devastating earthquakes in 2023 across various sections of the belt and neighboring regions, including the moment magnitude  $(M_W)$  7.8 and 7.6 Kahramanmaraş earthquake doublet in Türkiye and Syria (e.g., Mai et al., 2023; Ergintav et al., 2024), four  $M_W$ 6.3 earthquakes near Herat in Afghanistan (P. He et al., 2024), and the  $M_W$ 6.9 Al Haouz earthquake in Morocco (Yeck et al., 2023). In 2025, two additional large earthquakes struck the region: the  $M_W$ 7.1 Shigatse earthquake in southern Tibet and the  $M_W$ 7.7 Mandalay earthquake in Myanmar. Collectively, these events resulted in thousands of casualties and extensive economic losses, underscoring the urgent need for improved earthquake resilience and continental-scale studies of strain accumulation and release along the Alpides (Abdrakhmatov et al., 2024).

Previous efforts to compile GNSS velocities at regional and global scales have provided valuable insights into the deformation mechanisms along the Alpide belt (e.g., Hatzfeld & Molnar, 2010; Kreemer et al., 2014; Faccenna, Becker, Auer, et al., 2014; Devoti et al., 2017; Graham et al., 2018; Weiss et al., 2020; Serpelloni et al., 2022; Piña-Valdés et al., 2022; M. Wang & Shen, 2020; L. Wang & Barbot, 2023; Steffen et al., 2025; Y. Li et al., 2025). In addition, significant progress has been made in integrating interferometric synthetic aperture radar (InSAR) observations to illuminate areas with limited GNSS station coverage (e.g., Fattahi & Amelung, 2016; Barnhart, 2017; Hussain et al., 2018; Lazecký et al., 2020; Bletery et al., 2020; Weiss et al., 2020; Y. Li, Nocquet, Shan, & Song, 2021; Ou et al., 2022; Z. Liu & Wang, 2023; R. Jolivet et al., 2023; X. Huang et al., 2023a; Z. Huang et al., 2023; Watson et al., 2024; H. Chen et al., 2024; X. Li et al., 2024; Viltres et al., 2025). Some of these studies have specifically evaluated surface strain rates, enhancing our understanding of the spatial distribution and intensity of crustal deformation across different sections of the belt (e.g., Kreemer et al., 2014; Faccenna, Becker, Auer, et al., 2014; Weiss et al., 2020; M. Wang & Shen, 2020; Barbot & Weiss, 2021; Serpelloni et al., 2022; Piña-Valdés et al., 2022; L. Wang & Barbot, 2023; Z. Liu & Wang, 2023; Lindsey et al., 2023; H. Chen et al., 2024; Fang et al., 2024; Chousianitis et al., 2024; Steffen et al., 2025; Nucci et al., 2025; Y. Li et al., 2025).

Recent improvements in geodetic monitoring and station deployment have significantly increased the spatial density of GNSS networks across the Alpides, leading to new GNSS velocity fields for Anatolia (e.g., Kurt et al., 2023; Ergintav et al., 2023), the Euro-Mediterranean (e.g., Piña-Valdés et al., 2022; Serpelloni et al., 2022), northern Africa (e.g., Bougrine et al., 2019; Bahrouni et al., 2020; Billi et al., 2023), the Arabian Plate (Viltres et al., 2022), Tibet (e.g., Y. Li et al., 2025), and the India-Asia collision zone (e.g., M. Wang & Shen, 2020; Lindsey et al., 2023). Collectively, these studies provide a robust foundation for understanding regional tectonics and refining our estimates of strain rates, ultimately supporting seismic hazard assessment efforts along this extensive orogenic belt.

In this paper, we build upon those recent studies by compiling a unified, comprehensive, and up-to-date GNSS velocity field along the Alpine-Himalayan belt (hereafter referred to interchangeably as the Alpide belt or Alpides). Using this dataset, we estimate strain rates and posterior uncertainties for two focus regions: the eastern Mediterranean and the India-Asia collision zone, areas with the highest strain rates and dense GNSS station coverage. Our strain rate inversions explicitly account for velocity discontinuities across known creeping faults in the eastern Mediterranean, thereby avoiding the erroneous mapping of surface creep into strain rates associated with elastic coupling on faults, a distinction relevant for seismic hazard assessment.

Additionally, we demonstrate the utility of our combined GNSS velocity field for examining large-scale spatial patterns and relationships between GNSS-derived surface


strain rate styles and earthquake strain release styles from Kostrov-summed earthquake moment tensors (Dziewonski et al., 1981; Ekström et al., 2012), present-day stress release styles (Heidbach et al., 2025), and mantle dynamics inferred from seismic anisotropy compilations (Becker et al., 2012). Consequently, our study lays the foundation for future research on the links between strain accumulation and release, as well as the potential interplay between large-scale mantle dynamics and surface deformation along the Alpine-Himalayan belt.

# 2 Combined GNSS Velocity Field

In this study, we introduce FICORO\_GNSS (Castro-Perdomo, 2024), a Python software package developed to systematically filter and combine GNSS velocity fields. Utilizing this tool, we integrated 42 published velocity datasets across various sections of the Alpine-Himalayan belt (Table 1), resulting in a comprehensive GNSS velocity field comprising 11,177 horizontal and 3,940 vertical velocities (Fig. 2). Our methodology draws inspiration from the approaches of Piña-Valdés et al. (2022), who combined GNSS velocity solutions across Europe, and Zeng (2022), who developed the GNSS velocity field for the U.S. National Seismic Hazard Model. FICORO\_GNSS implements a robust filtering and combination process consisting of ten key steps:

- 1. Removing stations affected by postseismic transient motions: GNSS stations identified as being affected by postseismic transient motions in published velocity solutions were removed from the input data. This was accomplished by removing stations within a predefined radius around earthquake epicenters when previous studies explicitly reported postseismic transients. However, this approach represents only a first-order filter, as recent studies have shown measurable farfield deformation extending hundreds of kilometers from large earthquakes (e.g., Ergintav et al., 2024), and that postseismic transients may persist longer than previously assumed (S. Liu et al., 2020). Consequently, our combined velocity field and derived strain rate estimates might still include residual postseismic signals, particularly near the epicentral areas of the 2001 M<sub>s</sub>8.1 Kunlun earthquake in northern Tibet and the 2020 M<sub>W</sub>6.7 Elaziğ earthquake in eastern Türkiye, where some previous interseismic velocity fields inadvertently incorporated transient signals.
- 2. Filtering based on uncertainty distribution: We refined the input horizontal GNSS velocity fields by fitting a lognormal distribution to the east and north components of velocity uncertainties. Following the method outlined by Piña-Valdés et al. (2022), stations with uncertainties exceeding the 99th percentile of the fitted lognormal distribution were removed, as illustrated in Fig. S1.
- 3. Filtering based on spatial coherence of velocity magnitudes: We applied the standard (Z-Score) method used by Piña-Valdés et al. (2022) to identify and remove outlier GNSS stations based on velocity coherence with neighboring stations. GNSS stations whose velocity magnitudes in the East and North components deviated by more than two standard deviations,  $2\sigma$ , from the mean within a 20 km radius were removed. In tectonically active regions, a more lenient threshold of  $3\sigma$  was used to account for greater variability in GNSS velocities.
- 4. Aligning input GNSS velocity fields to ITRF2014: We applied a six-parameter Helmert transformation of GNSS velocities, solving for three translation rates  $\dot{\mathbf{T}} = (\dot{T}_x, \dot{T}_y, \dot{T}_z)$  and three rotation rates  $\dot{\mathbf{R}} = (\dot{R}_x, \dot{R}_y, \dot{R}_z)$ , fixing the scale-rate parameter  $(\dot{s})$  to zero. The translation- and rotation-rate vectors were estimated using a weighted least-squares inversion that minimizes horizontal velocity differences at common GNSS sites between the input and target velocity fields. The transformation of each input GNSS velocity field into the ITRF2014 (no-net-rotation) reference frame (Altamimi et al., 2017) is given by

$$\mathbf{V}_{\text{ITRF2014}} = \mathbf{V}_{\text{input}} + \dot{\mathbf{T}} + \dot{\mathbf{R}} \times \mathbf{X} + \dot{s} \, \mathbf{X},\tag{1}$$



Sea Fault, EAF: East Anatolian Fault, NAF: North Anatolian Fault, EAR: East African Rift, ATF: Altyn Tagh Fault, KF: Kunlun Fault, EHS: Eastern Himalayan Syntaxis, XJF: Xiaojiang Fault, XF: Xianshuihe Fault, JF: Jiali Fault, GYF: Ganzi-Yushu Fault, ZBSZ: Zhangjiakou-Bohai Seismic Zone, LJF: Lijiang Fault, HF: Ekström et al., 2012), with  $6 \ge M_W \ge 5$  earthquakes represented as dots. AEG: Aegean Sea Plate, WAEP: Western Anatolian Extensional Province, DSF: Dead al., 2024). Faults are sourced from AFEAD (Zelenin et al., 2021). Moment tensors for  $M_W \ge 6$  earthquakes are from the gCMT catalog (Dziewonski et al., 1981; Figure 1. Main tectonic features and seismicity along the Alpine-Himalayan belt. Slab depths are from Slab2 (Hayes et al., 2018) and ESHM2020 (Danciu et Haiyuan Fault, NQF: North Qilian Fault.

174

175

176

178

179

180

181

182

183

185

186

187

188

189

192

193

194

195

198

199

200

201

202

204

205

206

207

209

210

211

213

214

215

216

217

218

220

221

222

223

224

226

227

- where  $V_{\rm input}$  and  $V_{\rm ITRF2014}$  are the velocity vectors in the original and target reference frames, respectively, and  $\mathbf{X}=(X,Y,Z)$  is the station position vector (see Appendices A1-A2 in Altamimi et al., 2002). All input GNSS velocity fields listed in Table 1 were aligned to the velocity field of Serpelloni et al. (2022), expressed in the International GNSS Service (IGS) realization of the ITRF2014 reference frame (IGB14). Our implementation of this six-parameter Helmert transformation of GNSS velocities follows the same approach adopted in previous regional and global GNSS velocity combinations (e.g., Kreemer et al., 2006, 2009; Mallick et al., 2020; Graham et al., 2018; Gomez et al., 2020; Piña-Valdés et al., 2022; Serpelloni et al., 2022). The alignment statistics, including RMS alignment residuals and the number of common GNSS stations used for each transformation, are provided in Table 1. The consistently low RMS residuals demonstrate the quality of the transformation to ITRF2014 across all input GNSS velocity fields, with a mean RMS alignment residual of  $0.5 \pm 0.3$  mm yr<sup>-1</sup>.
- 5. Rotating input velocity fields to different reference frames: To aid visual inspection, tectonic interpretation, and subsequent manual filtering, we rotated input GNSS velocity fields, already aligned to ITRF2014, into various reference frames, including Eurasia, Nubia, India, East Anatolia, Arabia, Amuria, Yangtze (South China), Central Tibet, and Aegean-fixed reference frames, using the Euler poles listed in Table 2.
- 6. Filtering by velocity magnitude and azimuthal direction: We applied an Interquartile Range (IQR) method to assess the spread of velocity magnitudes and azimuthal directions at colocated stations where multiple velocities were reported by previous studies. Velocities falling below  $Q_1 1.5 \times IQR$  or above  $Q_3 + 1.5 \times IQR$  were considered outliers and subsequently removed, where  $Q_1$  and  $Q_3$  represent the first and third quartile, respectively. This approach was inspired by Zeng (2022), who filtered velocities that deviated by more than 20% in magnitude or azimuth from neighboring stations.
- 7. Combining velocity fields: Our approach for combining GNSS velocity fields builds on the method proposed by Zeng (2022), which averages velocity components for colocated stations and estimates uncertainties as the square root of the sum of averaged squared uncertainties from multiple studies. In our approach, instead of using the arithmetic mean, we computed the median velocities and uncertainties for colocated stations within a 0.01° (~1.1 km) radius, as the median is less sensitive to outliers. Fig. 2a shows the number of independent velocity observations used to compute the median horizontal velocity at each GNSS station. Vertical velocities were processed separately due to the limited availability of vertical rates in many previous studies, resulting in a less comprehensive dataset. To improve the vertical velocity field, we incorporated leveling data from Y. Wu et al. (2022) for the Tibetan Plateau and the India-Asia collision zone, which had already been aligned through a Helmert joint adjustment with regional continuous GNSS stations. To ensure robustness, we removed vertical velocities that deviated by more than  $2\sigma$  from the mean or exceeded the 99th percentile of the lognormal fit to the vertical uncertainty distribution, effectively filtering out outliers. However, vertical rates were not adjusted using a Helmert transformation and are provided as a separate dataset for completeness and future use. Vertical velocities are not used in our strain rate inversions, as we estimate the 2D strain rate tensor field solely from the combined horizontal GNSS velocities.
- 8. Manual filtering: We applied a manual filtering step to refine the combined velocity field, identifying outliers missed in previous filtering steps and stations affected by volcanic transient motions and anthropogenic sources. GNSS stations impacted by the Groningen gas field and volcanic activity at Campi Flegrei, Vesuvius, Etna, Afar, Santorini, and the Aeolian Islands were specifically excluded.
- 9. Scaling velocity uncertainties: We scaled horizontal velocity uncertainties in the combined velocity field using the method of Piña-Valdés et al. (2022), which

assumes GNSS velocity uncertainties follow a log-normal distribution. The uncertainties were adjusted to match the same percentile in a subjectively chosen target log-normal distribution. Here, we used the velocity uncertainty distribution of L. Wang and Barbot (2023), which has a mean close to 1 mm/yr for East and North velocity uncertainties, as shown in Fig. S2. This step was implemented to prevent unrealistically low velocity uncertainties, which could disproportionately weight certain stations and lead to improper scaling of the covariance matrix in our weighted least squares inversion for strain rates.

10. Final filtering based on uncertainty distribution: We removed stations with velocity uncertainties exceeding the 99th percentile of the scaled log-normal distribution, similar to our second filtering step but this time applied to the combined horizontal GNSS velocity field.

We intentionally included 2,714 additional horizontal GNSS velocities from stations located outside the Alpine-Himalayan belt to serve as tying sites for future global GNSS velocity combinations. This ensures that any future reference frame alignment involving our dataset can be performed robustly, as these stations provide broad geographic coverage for such transformations. Additionally, we provide rotated velocity fields in ten different reference frames, openly available in a public repository (Castro-Perdomo, 2025), to further facilitate their use in future research. Corresponding maps of the rotated horizontal GNSS velocity fields are included in the supplementary material (Figs. S3-S5).

#### 3 Strain Rate Inversion

Given the spatially scattered nature of GNSS velocities, interpolating these data is essential for estimating surface strain rates. Various methodologies have been developed for this purpose, ranging from calculating strain rates using baselines—often through techniques like Delaunay triangulation (e.g., J. C. Savage, 1983; J. Savage et al., 2001; Cai & Grafarend, 2007; Kreemer et al., 2018, 2020)—to employing non-parametric and elastic basis functions to fit surface velocities (e.g., A. Haines & Holt, 1993; Beavan & Haines, 2001; Tape et al., 2009; A. J. Haines et al., 2015; Sandwell & Wessel, 2016; Okazaki et al., 2021; Johnson, 2024). Additionally, methodologies utilizing weighted mean and locally-weighted regression have been explored (e.g., El-Fiky & Kato, 1998; Z.-K. Shen et al., 1996, 2015), and recent studies have proposed Bayesian interpolation schemes (e.g., Pagani et al., 2021) and geostatistical approaches based on ordinary kriging (e.g., Maurer & Materna, 2023).

Here, we adopted the body-force method proposed by Johnson (2024), as implemented in the BForStrain MATLAB code (Johnson, 2023). BForStrain interpolates 2D velocity fields using analytical Green's functions for a thin elastic body under in-plane body forces (Sandwell & Wessel, 2016). Velocities and strain rates are estimated at the centroids of triangles within a 2D adaptive triangular mesh generated using MESH2D, a Delaunay mesh-generator by Engwirda (2009), whose resolution is refined based on the GNSS station density. BForStrain enables uncertainty estimation through linear error propagation of GNSS velocity uncertainties and posterior sampling across a range of regularization parameters. Notably, it also allows for velocity discontinuities across predefined creeping faults, which are conflated with strain in most of the other methods.

Our strain rate inversions solve for the distribution of body forces at the vertices of a triangular mesh and surface creep rates on faults that best fit the observed geodetic surface velocities  $v_{\rm obs}$ . The regularized weighted least squares formulation can be expressed as:

Table 1. GNSS velocity fields used in the combination, and their alignment statistics.

Source	Study area	Sites	Reference frame	Ties	RMS (mm/yr)
Alchalbi et al. (2010)	Northern Dead Sea Fault	62	ITRF2000	14	0.45
Bahrouni et al. (2020)	Tunisia	42	ITRF2014	38	0.26
Billi et al. (2023)	Northern Africa	431	Eurasia ITRF2014	292	0.26
Bougrine et al. (2019)	Algeria	313	ITRF2014	259	0.37
Briole et al. (2021) <sup>⊕</sup>	Aegean	329	ITRF2014	188	0.97
Brockmann et al. (2019) <sup>⊕</sup> ®	Euromediterranean (EPN)	3047	Eurasia	2492	0.16
Brockmann et al. (2019) <sup>⊕</sup> ®	Switzerland (EPN-CH8)	214	Eurasia	245	1.63
Castro-Perdomo et al. (2022) <sup>⊕</sup>	Southern Dead Sea Fault	66	ITRF2014	56	0.33
England et al. (2016) <sup>®</sup>	Anatolia	346	Eurasia	26	0.68
Ergintav et al. (2023) <sup>⊕</sup>	Anatolia	732	Anatolia ITRF2014	108	1.04
Floyd et al. (2023) <sup>⊕</sup>	Hellenic subduction zone	360	Eurasia	70	1.34
Gomez et al. (2020) <sup>⋄</sup>	Dead Sea Fault	181	ITRF2008	26	1.15
Graham et al. (2018) <sup>®</sup>	Global	8354	North America	1957	0.50
Hamiel and Piatibratova (2021) <sup>⋄</sup>	Dead Sea Fault	291	ITRF2014	57	0.90
R. Jolivet et al. (2023) <sup>⊕</sup>	North Anatolian Fault	81	ITRF2014	51	0.46
Kadirov et al. $(2014)^{\odot}$	Caucasus - Azerbaijan	110	Eurasia ITRF2000	50	0.26
Karakhanyan et al. (2013) <sup>⋄</sup>	Armenia and Lesser Caucasus	55	Eurasia ITRF2008	49	0.26
Khorrami et al. (2019) <sup>⋄</sup>	Iran	399	Eurasia ITRF2008	21	0.45
Kurt et al. (2023)	Anatolia	836	Eurasia	8	0.57
Y. Li et al. (2025) <sup>®</sup>	Tibet-SE Asia	4458	Eurasia-fixed	101	0.59
Liang et al. $(2013)^{\oplus}$	Tibetan Plateau	750	Eurasia ITRF2008	18	1.07
McClusky et al. (2010)	Afar triple junction	58	Nubia	19	0.54
Nocquet (2012) <sup>®</sup>	Euromediterranean	1495	Eurasia	757	0.36
Özarpacı et al. (2021)	North Anatolian Fault	61	Eurasia	23	0.25
Özbey et al. (2024) <sup>\dagger</sup>	Cyprus and East Anatolia	98	ITRF2014	20	0.21
Özdemir and Karshoğlu (2019) <sup>\dagger</sup>	Anatolia	213	Eurasia	41	0.25
Özkan et al. (2023) <sup>\dagger</sup>	Hatay Triple Junction	65	Eurasia	40	0.26
Perry et al. (2019)	Pamir-Hindu Kush	82	Eurasia ITRF2008	23	0.62
Piña-Valdés et al. (2022) <sup>⊕</sup> ®	Euromediterranean	4837	ITRF2014	2871	0.37
Reilinger et al. (2006)	Euromediterranean	433	Eurasia	178	0.72
Saleh and Becker (2014)	Egypt	149	Eurasia	122	0.61
Serpelloni et al. (2022) <sup>⊕</sup>	Euromediterranean	3350	ITRF2014	*	*
Sokhadze et al. (2018) <sup>⊕</sup>	Caucasus - Georgia	72	Eurasia	24	0.21
Stamps et al. $(2018)^{\diamond}$	East African Rift	153	Nubia	95	0.48
Tatar et al. (2012)	North Anatolian Fault	48	Eurasia	30	0.56
Viltres et al. (2020)	Danakil - Southern Red Sea	33	Nubia	4	0.13
Viltres et al. (2022) <sup>⊕</sup>	Arabia	189	ITRF2014	27	0.33
L. Wang and Barbot (2023) <sup>®</sup>	India-Eurasia collision	2870	Local NNRT <sup>⊙</sup>	55	0.53
M. Wang and Shen (2020) <sup>⋄</sup>	China	3273	China	99	0.49
Wedmore et al. $(2021)^{\diamond}$	Africa	262	ITRF2014	150	0.44
Zubovich et al. (2016)	Pamir and Tian Shan	411	Eurasia	22	0.62
Zheng et al. (2017) <sup>\display</sup>	India-Eurasia collision	2576	Eurasia	46	0.48

<sup>\*</sup> Reference velocity field into which all others are aligned.

Table 2. Euler poles used to rotate the combined horizontal GNSS velocity field.

Plate	$\omega_x$ (° $Myr^{-1}$ )	$\omega_y$ (° $Myr^{-1}$ )	$\omega_z$ (°Myr <sup>-1</sup> )	Lat	Lon	$\omega$ (°Myr <sup>-1</sup> )	Source	Map
Eurasia*	-0.023500	-0.147600	0.214000	55.07	-99.05	0.26102	Altamimi et al. (2017)	Fig. 2b
Arabia*	0.328400	-0.035040	0.406820	50.93	-6.09	0.52400	Viltres et al. (2022)	Fig. S3a
Nubia*	0.027400	-0.170400	0.203700	49.73	-80.87	0.26698	Altamimi et al. (2017)	Fig. S3b
East Anatolia*	1.008722	0.543127	1.020384	1.01	0.54	1.02038	Ergintav et al. (2023)	Fig. S3c
West Aegean <sup>⋆</sup>	0.050898	0.147426	0.158303	45.43	70.95	0.22223	Ergintav et al. (2023)	Fig. S4a
Sinai*	0.242500	-0.047970	0.349390	54.72	-11.19	0.42800	Hamiel and Piatibratova (2021)	Fig. S4b
India*	0.320500	-0.001400	0.403800	51.56	-0.25	0.51554	Altamimi et al. (2017)	Fig. S4c
Amuria*	-0.048000	-0.107140	0.274140	66.82	-114.13	0.29822	This study <sup>©</sup>	Fig. S5a
South China*	-0.057800	-0.127200	0.293300	64.53	-114.45	0.32490	Vardić et al. (2022)	Fig. S5b
Central Tibet*	0.129830	-0.713600	-0.023360	-1.84	-79.69	0.72569	This study <sup>®</sup>	Fig. S5c

<sup>★</sup> Relative to the International Terrestrial Reference Frame 2014 (ITRF2014, Altamimi et al., 2017)

 $<sup>\</sup>circledast$  GNSS velocity fields that are themselves combinations of previous datasets (see Table S2).


 $<sup>\</sup>diamond$  GNSS velocity fields that incorporate data from earlier studies to densify GNSS station coverage.

<sup>⊙</sup> First rotated into Karakhanyan et al. (2013) velocity field.

 $<sup>\</sup>oplus$  GNSS velocity fields reporting vertical velocities.

<sup>⊙</sup> No-Net-Rotation and Translation.

 $<sup>\</sup>odot$  Estimated using three GNSS stations within the Amurian plate (see Fig. S6a,b for details).



**Figure 2.** (a) Number of independent horizontal velocity solutions at each GNSS station. (b) Combined horizontal GNSS velocity field expressed in a Eurasia-fixed reference frame. (c) Combined vertical GNSS velocity field (red: uplift, blue: subsidence).

$$\begin{bmatrix} \Sigma_v^{-\frac{1}{2}} v_{\text{obs}} \\ \alpha c \\ 0 \end{bmatrix} = \begin{bmatrix} \Sigma_v^{-\frac{1}{2}} G_v & \Sigma_v^{-\frac{1}{2}} G_c \\ 0 & \alpha I \\ \beta I & 0 \end{bmatrix} \begin{bmatrix} f \\ s_c \end{bmatrix}, \tag{2}$$

where  $\Sigma_v$  represents the covariance matrix of the observed velocities, I is the identity matrix, and  $G_v$  and  $G_c$  are matrices of Green's functions for body forces and fault creep rates, respectively. The objective is to solve for the body forces and fault creep rates (f and  $s_c$ ) that minimize the residual between observed and predicted surface velocities, subject to Tikhonov regularization constraints ( $\alpha$  and  $\beta$ ) on surface creep rates and body-force magnitudes, respectively.

Upon solving the linear system in Eq. 2, we aim to calculate the full posterior probability distribution for the estimated strain rates. This involves computing the predicted strain rates,  $\varepsilon_{\beta}$ , from the best-fitting model of body forces and surface creep rates and then deriving their covariance,  $\Sigma_{\varepsilon,\beta}$ , to quantify uncertainties. The posterior probability distribution for the strain rates, given the observed data d and regularization parameter  $\beta$ , can be expressed as:

$$p(\varepsilon|d,\beta) = \frac{1}{\left[(2\pi)^k |\Sigma_{\varepsilon,\beta}|\right]^{\frac{1}{2}}} \exp\left(-\frac{1}{2}(\varepsilon - \varepsilon_\beta)^\top \Sigma_{\varepsilon,\beta}^{-1}(\varepsilon - \varepsilon_\beta)\right). \tag{3}$$

To encapsulate uncertainty in the body force damping  $\beta$ , we approximated the posterior distribution  $p(\varepsilon|d)$  by drawing 500 random samples from Eq. 3 for each  $\beta$  value, representing altogether discrete realizations of the posterior distribution  $p(\varepsilon|d)$ , following the same procedure implemented by Johnson (2024). We then evaluated the solution's smoothness and fit across a range of damping parameters, with L-curves provided in the supplementary material to support our subjective damping choices (Fig. S7). The damping parameter  $\alpha$  controlling surface creep rates was selected subjectively to ensure that the model neither overfits nor underfits observed GNSS velocities across well instrumented creeping faults. Finally, strain rates are computed from the inverted model solution of body forces and surface creep rates (f and  $s_c$ ) and analytical Green's functions ( $G_{\varepsilon}, G_{c,\varepsilon}$ ) relating those to strain rates, for a particular  $\beta$  value:

$$\varepsilon_{\beta} = G_{\varepsilon} f_{\beta} + G_{c,\varepsilon} s_{c\beta}. \tag{4}$$

DOI: 10.1029/2025JB031738

We conducted three strain rate inversions. The first one aimed to capture large-scale deformation patterns and identify regions with the highest deformation rates along the entire Alpine-Himalayan belt (Fig. 3). This inversion applied moderate body-force damping constraints to emphasize broad-scale strain rate features, without estimating posterior uncertainties or surface creep rates. Based on these results, we selected two regions for detailed analysis: the eastern Mediterranean and the India-Asia collision zone, both characterized by high strain rates and dense GNSS station coverage. For these regions, we performed focused inversions using refined triangular meshes to resolve finer-scale deformation features, as discussed in Section 4.

In these focused inversions, we estimated posterior uncertainties by drawing 3,000 samples from the posterior distribution, utilizing six damping parameters near the L-curve elbows to balance model smoothing and fit to observed GNSS velocities. To further regularize our least squares inversion, we incorporated priors on fault creep rates in the eastern Mediterranean based on creep rate estimates from previous geodetic studies along select fault segments. Specifically, we included shallow creep rate priors from the Izmit and İsmetpaşa segments of the North Anatolian fault (Cakir et al., 2005; Kaneko et al., 2013; Cetin et al., 2014; Hussain et al., 2016; Aslan et al., 2019; Özarpacı et al.,

2021; Kurt et al., 2023), the Marmara segment of the North Anatolian Fault (Yamamoto et al., 2019), the Palu segment of the East Anatolian fault (Cavalié & Jónsson, 2014; Bletery et al., 2020; Cakir et al., 2023), and the northern segment of the Jordan Valley fault located south of the Sea of Galilee, where shallow fault creep has been reported (Hamiel et al., 2016; Hamiel & Piatibratova, 2021).

Considering that the method proposed by Johnson (2024) estimates strain rates in a Cartesian coordinate system, evaluating distortion effects across the continental-scale Alpine-Himalayan belt is crucial. To this end, we generated a synthetic Euler-pole-predicted velocity field over the entire belt using an equidistant grid with a  $1^{\circ} \times 1^{\circ}$  spacing and inverted this synthetic velocity field to estimate strain rates, applying the same body force damping parameters used in our large-scale inversion (Fig. S8). The purpose of this test is to isolate distortion artifacts: since the synthetic Euler-pole-predicted velocity field represents purely rigid body rotations, any nonzero strain rates estimated far from tectonic plate boundaries can be attributed to coordinate system distortions rather than true deformation.

Our analysis revealed that strain rates resulting from distortion effects are small, remaining below 5 nanostrain/yr. Faster-moving plates such as India and Arabia exhibit the largest distortion effects, which increase with the angular distance from each plate's Euler pole. Additionally, distortion more significantly impacts the magnitude of dilatation rates compared to the second invariant and maximum shear strain rates. Furthermore, distortion effects are spatially systematic and intensify with latitude, becoming more pronounced near the geographic poles, a factor not highlighted in our test due to our use of a Eurasia-fixed reference frame for the GNSS velocities. Further details on our test for distortion effects are provided in Fig. S8 in the supplementary material.

Several scalar measures can be used to characterize the strain rate tensor, often expressed in terms of its invariants or eigenvalues (see Appendix A). In this study, we adopt two widely used scalar quantities: the first invariant  $(I_1^{2D}, \text{ dilatation rate})$  and the Frobenius norm of the 2D horizontal strain rate tensor  $(\|\dot{\varepsilon}\|_F^{2D})$ , defined as:

$$I_1^{2D}(\dot{\varepsilon}) = \dot{\Delta} = \operatorname{tr}(\dot{\varepsilon}) = \dot{\varepsilon}_{11} + \dot{\varepsilon}_{22} = \dot{\varepsilon}_1 + \dot{\varepsilon}_2, \tag{5}$$

$$I_1^{2D}(\dot{\varepsilon}) = \dot{\Delta} = \text{tr}(\dot{\varepsilon}) = \dot{\varepsilon}_{11} + \dot{\varepsilon}_{22} = \dot{\varepsilon}_1 + \dot{\varepsilon}_2,$$

$$\|\dot{\varepsilon}\|_F^{2D} = \sqrt{\sum_{i,j} \dot{\varepsilon}_{ij}^2} = \sqrt{\dot{\varepsilon} : \dot{\varepsilon}} = \sqrt{\text{tr}(\dot{\varepsilon}^2)} = \sqrt{\dot{\varepsilon}_{11}^2 + \dot{\varepsilon}_{22}^2 + 2\dot{\varepsilon}_{12}^2}.$$
(6)

For clarity, throughout this paper we refer to the Frobenius norm of the 2D horizontal strain rate tensor as "strain rate" (e.g., Tape et al., 2009). Note that for the deviatoric strain rate tensor  $\dot{\varepsilon}'$ , the Frobenius norm is directly related to the deviatoric second invariant,  $\|\dot{\varepsilon}'\|_F = \sqrt{2J_2}$ , which in two dimensions is equivalent to  $\|\dot{\varepsilon}'\|_F^{2D} = \sqrt{2}\,\dot{\gamma}_{\rm max}^{2D}$ , where  $\dot{\gamma}_{\max}^{2D} = \sqrt{J_2} = \frac{1}{2}(\dot{\varepsilon}_1 - \dot{\varepsilon}_2)$  denotes the maximum shear strain rate.

# 4 Velocity and Strain Rate Fields

316

317

318

320

321

322

325

326

327

329

330

331

332

333

334

337

338

339

341

342 343

345

346 347

349

350

351

352

353

354

355

359

360

## 4.1 Combined Horizontal GNSS Velocity Field

The horizontal velocity field relative to Eurasia (Fig. 2b) reveals coherent largescale motions that underscore the region's active tectonics and geodynamics, which have been extensively studied over the past 50 years (e.g., Molnar & Tapponnier, 1975; D. McKenzie, 1978; Jackson & McKenzie, 1984; Şengör, 1986; McClusky et al., 2000; Reilinger et al., 2006; Becker & Faccenna, 2011; Nocquet, 2012; L. Jolivet et al., 2013; Hatzfeld & Molnar, 2010; Kreemer et al., 2014; Palano, 2014; Faccenna, Becker, Auer, et al., 2014; L. Jolivet et al., 2018; Barbot & Weiss, 2021; M. Wang & Shen, 2020; L. Wang & Barbot, 2023). Notable large-scale features include the counterclockwise rotation of Arabia, western Iran, Anatolia, and the Aegean (e.g., McClusky et al., 2000; Reilinger et al., 2006); the acceleration of Anatolia toward the Hellenic Trench, likely driven by trench retreat and slab rollback (e.g., Le Pichon & Angelier, 1979; Royden, 1993; Reilinger et al., 2006;

Faccenna, Becker, Auer, et al., 2014); and lateral escape tectonics on both flanks of the Tibetan Plateau as a result of the India-Asia collision (Tapponnier et al., 1982; M. Wang & Shen, 2020).

Additionally, the horizontal velocity field in a Eurasia-fixed reference frame reveals a clockwise rotation around the eastern Himalayan syntaxis in Southeast Asia (e.g., Tapponnier et al., 1982; Holt et al., 1991; Yin & Harrison, 2000; Sol et al., 2007; Hatzfeld & Molnar, 2010; M. Wang & Shen, 2020). Several mechanisms have been invoked to explain deformation patterns observed in this complex region, including crustal and mantle flow driven by gravitational potential energy (GPE) release from the Tibetan Plateau (e.g., Molnar & Lyon-Caen, 1988; Royden et al., 1997; F. Shen et al., 2001; Bai et al., 2010; X. He et al., 2021; Timsina et al., 2024), localized flow due to partial melting (Sun et al., 2025), and the possible influence of slab rollback (e.g., F. Shen et al., 2001; León Soto et al., 2012; Hou et al., 2024).

## 4.2 Combined Vertical GNSS Velocity Field

The vertical velocity field illustrated in Fig. 2c also displays coherent regional and large-scale tectonic and geodynamic signatures. Notable features include the well-documented uplift in the Fennoscandian Peninsula, with vertical rates reaching ~10 mm/yr, attributed to glacial isostatic adjustment (GIA) following the Late Pleistocene glaciation (e.g., Milne et al., 2001; Johansson et al., 2002; Nocquet, 2012). Additionally, ~1 mm/yr uplift rates are observed in the Eifel volcanic field in Germany, linked to a buoyant mantle plume (Ritter et al., 2001; Walker et al., 2005; Kreemer et al., 2020), and 1-2.5 mm/yr uplift rates are recorded along the Alps (e.g., Nocquet et al., 2016; Serpelloni et al., 2022; Piña-Valdés et al., 2022). These uplift rates are interpreted to result from a mix of processes, with contributions from shortening (e.g., van Gelder et al., 2017), erosion (e.g., Kuhlemann et al., 2002), isostatic rebound after the last deglaciation (e.g., Mey et al., 2016), slab detachment (e.g., Sue et al., 1999; Wortel & Spakman, 2000; Fox et al., 2015), and Eurasia-Adria subduction (e.g., Lippitsch et al., 2003; Brückl et al., 2007), as discussed by Sternai et al. (2019).

Further features of the vertical velocity field include widespread subsidence in the Po Basin, with rates exceeding 5 mm/yr. This subsidence is driven by a combination of anthropogenic sources (e.g., groundwater extraction), Adriatic slab retreat, and compaction of alluvial deposits (e.g., Carminati et al., 2003; Tosi et al., 2013; Serpelloni et al., 2013). In contrast, the central Apennines exhibit both uplift and positive dilatation rates (see Fig. S9), consistent with active mantle flow in a slab gap and recent orogeny (Shaw & Pysklywec, 2007; Serpelloni et al., 2013; Faccenna, Becker, Miller, et al., 2014).

Systematic uplift is also evident along the Levant, where vertical rates reach up to ~2 mm/yr (e.g., Serpelloni et al., 2013, 2022), associated with the activity of the Dead Sea Fault (e.g., Ten Brink et al., 1990; Gomez et al., 2006, 2020). Finally, the vertical velocity field highlights both widespread uplift across the Tibetan Plateau and subsidence along its southeastern flank around the eastern Himalayan syntaxis (Y. Wu et al., 2022). Observed uplift rates ranging from ~1 to ~4 mm/yr have been attributed to vertical crustal thickening driven by the India-Asia collision (e.g., England & Houseman, 1986; Molnar et al., 1993; S. Liu et al., 2025), while the 0.5-3.5 mm/yr subsidence rates in the southeast have been interpreted as a consequence of crustal flow and gravitational collapse (Y. Wu et al., 2022).

## 4.3 Strain Rate Field

Continental-scale strain rate maps and profiles across sections of the Alpide belt are presented in Figs. 3 and 4. To enhance the readability of our strain rate maps, we regridded the strain rate tensor components from the original triangular meshes to an

equidistant grid with  $0.1^{\circ} \times 0.1^{\circ}$  spacing using linear interpolation. Regional-scale strain rate features derived from our combined velocity field align with previous studies (e.g., Kreemer et al., 2014; Weiss et al., 2020; M. Wang & Shen, 2020; Serpelloni et al., 2022; Piña-Valdés et al., 2022; L. Wang & Barbot, 2023; Lindsey et al., 2023; Y. Li et al., 2025), although differences in strain rate magnitudes are observed. These discrepancies are primarily due to the significant recent expansion of GNSS networks—particularly in the eastern Mediterranean, northern Africa and the India-Asia collision zone—and the inherent non-uniqueness of strain rate estimates. The increased GNSS station density in these regions has substantially improved resolution compared to earlier GNSS-derived strain rate models.

Below, we discuss key features observed in the velocity and strain rate fields for the eastern eastern Mediterranean and the India-Asia collision zone. These regions were selected due to their dense GNSS station coverage, which has improved significantly with the addition of over 2,000 new stations in recent years (M. Wang & Shen, 2020; Kurt et al., 2023; Y. Li et al., 2025), and the presence of high strain rate belts, such as those in Anatolia, Tibet, Himalayas, and the southeast borderland of the Tibetan Plateau. For these areas, we performed additional strain rate inversions using finer triangular meshes, with a nominal spacing of 100 km in regions with sparse GNSS coverage, refined to approximately 5 km in areas with dense instrumentation. Unlike our large-scale inversion, where we prioritized regional trends by subjectively selecting a range of body-force damping parameters  $(\beta)$  that produce a smooth strain rate field, these focused inversions applied an L-curve method to identify a range of damping parameters within the elbow of the L-curve, optimizing the balance between model roughness and fit to the data. Consequently, the refined strain rate estimates presented below have higher resolution and exhibit greater magnitudes in highly deforming regions compared to those from our broaderscale inversion. Moreover, unlike our large-scale inversion, we compute and display posterior uncertainties for the refined strain rate estimates, as shown in Fig. S14 of the supplementary material.

# 4.3.1 Eastern Mediterranean

Fig. 5 presents the results of our refined strain rate inversion for the eastern Mediterranean region. Panel (a) displays the GNSS velocity field in the East Anatolia-fixed reference frame from Ergintav et al. (2023), highlighting the pronounced N-S extension in western Anatolia—a feature that is less evident in the commonly used Eurasia-fixed reference frame. This reference frame also emphasizes the right-lateral North Anatolian Fault, where slip rates range from ~20 to ~30 mm/yr, increasing westward (e.g., McClusky et al., 2000; Reilinger et al., 2006; Le Pichon & Kreemer, 2010; Hussain et al., 2018; Kurt et al., 2023; Ergintav et al., 2023). Additionally, the velocity field reveals <10 mm/yr of left-lateral motion between the Anatolian and Arabian plates along the East Anatolian Fault (e.g., Güvercin et al., 2022), with negligible fault-perpendicular velocities. This observation, first noted by Reilinger et al. (2006), was cited as evidence against the extrusion hypothesis as the primary driver of Anatolia's motion (Şengör et al., 1985), instead supporting an alternative model in which deformation is primarily controlled by slab rollback and trench retreat (e.g., Le Pichon & Angelier, 1979; Malinverno & Ryan, 1986; D. McKenzie & Yılmaz, 1991; Royden, 1993; Heidbach, 2005; Faccenna et al., 2006).

Panel (b) in Fig. 5 illustrates the triangular mesh used in the refined strain rate inversion, with higher resolution applied in densely instrumented areas and along creeping fault segments. Well-instrumented regions include most of the Anatolian Plate, the Peloponnese, Crete, northern Cyprus, and the Levant. In contrast, areas with sparse GNSS coverage include Libya in north Africa, the northern Arabian Plate across Syria and Iraq, and the region north of the Greater Caucasus near the Caspian Sea. Panel (c) presents a map of dilatation rates and principal strain rate directions, illustrating a transition from E-W extension in central Anatolia to N-S extension across the western Anatolia exten-





Figure 3. (a) Triangular mesh used in the large-scale strain rate inversion, with a nominal node spacing of 100 km in regions with sparse GNSS coverage (warmer colors) and refined to  $\sim 15$  km in well-instrumented areas (cooler colors). The locations of four profiles shown in Fig. 4, are also indicated. (b) First invariant of the strain rate tensor (dilatation rate) with principal strain rate directions plotted on a  $0.75^{\circ}$  equidistant grid. Red indicates extension, while blue represents contraction. (c) Strain rate, with a lower threshold of 10 nanostrain/yr to emphasize the main deformation belts and shear zones. We applied a mask to suppress strain rate bleeding artifacts in poorly instrumented regions and over water bodies, including the Mediterranean Sea, western Red Sea borderlands, Indian Ocean, Bay of Bengal, and South China Sea (see Fig. S10).




and elevation. Profile A-A' spans from the Adriatic Sea to the Caspian Sea. Profile B-B' extends from Cyprus to the Black Sea. Profile C-C' traverses from the Tian Figs. S11, S12 and S13 in the supplementary material. KTJ: Karliova Triple Junction, EAP: Eastern Anatolia Plateau, NAF: North Anatolian Fault, KKF: Karako Profiles of GNSS velocities in a Eurasia-fixed reference frame, strain rates, and topography across various sections of the Alpine-Himalayan belt. Pro-Shan to Southeast Asia. Profile D-D' runs from the Indo-Gangetic Plain to the Baikal Rift. Additional profiles along other sections of the Alpide belt are shown in file locations are shown in Fig. 3a. From top to bottom, the rows display along-profile velocity, across-profile velocity, vertical velocity, strain rate, dilatation rate, rum Fault, MFT: Main Frontal Trust, KF: Kunlun Fault, ATF: Altyn Tagh Fault. Figure 4.

sional province. Principal strain directions also delineate a shear belt extending offshore from the North Anatolian Fault into the North Aegean Sea along the North Aegean Trough (e.g., Flerit et al., 2004; Rodriguez et al., 2023), which widens westward, consistent with recent results by Chousianitis et al. (2024), as shown in Fig. 5c, d. The southwestern Aegean exhibits low strain rates (< 10 nanostrain/yr), supporting previous studies suggesting that portions of the southwestern Aegean Sea behave as a relatively rigid block (e.g., McClusky et al., 2000; Kreemer & Chamot-Rooke, 2004). Additionally, mild extension (10–20 nanostrain/yr) is observed across parts of the south central Taurus mountains, as depicted in profile B-B' in Fig. 4.

Although GNSS station coverage in the Caucasus region is sparser than in Anatolia, our results reveal a west-to-east transition from predominantly N-S to NNE-SSW shortening across the Greater Caucasus, with strain rates increasing toward the Caspian Sea, exceeding 60 nanostrain/yr. In contrast, the Lesser Caucasus exhibits localized, discontinuous zones of extension and shortening, with principal strain directions indicating a combination of strike-slip motion, NW-SE shortening, and NE-SW extension (see Fig. 5c). These findings align with geological studies documenting the complex coexistence of compressional and extensional structures (e.g., Rebaï et al., 1993) and evidence of syn- and post-collisional magmatism (Ismail-Zadeh et al., 2020, and references therein). Similarly, our results agree with early block model interpretations by Reilinger et al. (2006), which, despite being based on a more limited GNSS velocity dataset, inferred small fault-normal extension within the Lesser Caucasus and south of the Caspian Sea (see their Figs. 9b, 9c, and 11). However, a denser GNSS network is needed to confirm these observations.

Panel (d) in Fig. 5 highlights strain rates in Anatolia, revealing along-strike variations in strain rates along the NAF. These variations align with findings by Z. Liu and Wang (2023), who reported a potential link between elevated strain rates, surface creep, and earthquake ruptures over the past century, starting with the 1939  $M_W7.8$  Erzincan Earthquake. Along-strike strain rate variations were also noted by Weiss et al. (2020). although they emphasized biases due to sparse GNSS station coverage prior to the recent expansion of the GNSS network in Türkiye (Kurt et al., 2023; Ergintav et al., 2023). If these variations reflect afterslip-related surface creep, they suggest that interseismic velocity fields in Anatolia could be influenced by long-term postseismic afterslip transients, as documented along the Izmit rupture 20 years after the 1999 earthquake (Ozarpacı et al., 2021). Furthermore, InSAR data and stress-driven afterslip models suggest that postseismic afterslip along the creeping segment of the East Anatolian Fault (EAF) could persist for several decades (>50 years) following the 2023 Kahramanmaraş earthquake doublet (Z. Liu et al., 2025). However, the origin and time-dependent evolution of surface creep along various segments of the NAF and EAF remain debated (e.g., Cakir et al., 2012; Kaneko et al., 2013; Cetin et al., 2014; Bilham et al., 2016; R. Jolivet et al., 2023; Z. Liu et al., 2025).

The study by Özarpacı et al. (2021) also reported that current locking depths along the Izmit segment of the NAF are significantly shallower than pre-earthquake estimates, attributing this reduction to afterslip. However, given that transient signals observed in GNSS velocity profiles extend up to 70 km from the fault, viscoelastic mantle relaxation cannot be ruled out as a contributing factor, as proposed by earlier studies (e.g., L. Wang et al., 2009; Hearn et al., 2009; Ergintav et al., 2009; DeVries et al., 2017). If ongoing post-seismic viscoelastic relaxation is confirmed, it would challenge the hypothesis by Hussain et al. (2018), which proposed a short-lived postseismic period of approximately 10 years. The discrepancy may arise from Hussain et al. conflating two distinct parameters: the locking depth (16 km) derived from elastic half-space modelling of surface velocities (e.g., J. C. Savage & Burford, 1973) and the elastic thickness assumed in layered viscoelastic models (e.g., J. Savage & Prescott, 1978). Further research incorporating vertical GNSS and InSAR velocities (e.g., Jónsson, 2008; Fukuda & Johnson, 2021; Luo & Wang, 2022) is necessary to better distinguish between the contributions of afterslip and viscoelas-



where the strain rate is larger than 10 nanostrain/yr. (d) Strain rate. The panel in the bottom right compares modeled strain rates from inversions that accounted Dilatation rate (first invariant) and principal strain rate directions (red: stretching, blue: shortening). Principal strain rate directions are shown only at locations Fault, as indicated by the white box on the map. PEL: Peloponnese, GC: Gulf of Corinth, WAEP: Western Anatolian Extensional Province, NAT: North Aegean (a) GNSS velocity field rotated into the East Anatolia-fixed reference frame from Ergintav et al. (2023), highlighting the transition from E-W exred dots) and ignored (dark gray dots) the contribution of surface creep to strain rates along a profile crossing the İsmetpaşa segment of the North Anatolian tension in central Anatolia to N-S extension in western Anatolia. (b) Triangular mesh used for strain rate inversion, color-coded by normalized triangle area. Frough, NAF: North Anatolian Fault, EAF: East Anatolian Fault, KTJ: Karliova Triple Junction, EAP: Eastern Anatolian Plateau. Figure 5.

tic mantle flow along the North Anatolian Fault, which is thought to be underlain by a 50–100 km-wide shear zone in the lower crust and upper mantle (e.g., Taylor et al., 2016; Papaleo et al., 2018; Barbot & Weiss, 2021).

Our results also reveal a shear belt along the East Anatolian Fault (EAF), though strain rates are notably lower than those along the North Anatolian Fault (NAF), typically ranging between 50 and 80 nanostrain/yr, in agreement with previous InSAR- and GNSS-derived strain rate estimates (Weiss et al., 2020; Özbey et al., 2024). Localized elevated strain rates are observed near the epicenter of the 2020  $M_W$  6.7 Elaziğ earthquake, as also pointed out by Weiss et al. (2020), likely reflecting postseismic transients inadvertently incorporated into prior interseismic velocity fields. Another area of elevated strain rates is the Karlıova triple junction, where strain rates exceed 100 nanostrain/yr. Additionally, increased posterior uncertainties are evident along the main shear belts of the NAF, EAF, DSF, and the Gulf of Corinth, as detailed in Fig. S14 in the supplementary material.

To assess the quality of our GNSS velocity inversion for strain rates, we compared observed and model-predicted (i.e., mean posterior) GNSS velocities, as shown in Fig. 6a. The results indicate excellent agreement, with 80% of residual velocities falling within  $\pm 1 \text{ mm/yr}$ , as illustrated in the histogram in the bottom right of Fig. 6b. We further quantified the spatial coherence in the residual velocities by computing a "systematic misfit index" (e.g., Johnson et al., 2024), defined as the absolute value of the average dot product of normalized residual velocity vectors with their neighbors within a specified radius (~50 km), which reveals no evidence for regional-scale systematic residuals (Fig. 6b). Our model also effectively resolves velocity gradients across closely spaced creeping and locked fault segments where GNSS station density is high. For example, panels 1 and 2 in Fig. 6a show that it accurately captures the velocity gradients across the creeping Ismetpaşa segment and the locked Düzce segment of the NAF, separated by 100 km. Panels 1 and 2 in Fig. 6a further demonstrate strong correspondence between observed, mean posterior, and predicted fault-parallel velocities from an elastic half-space antiplane dislocation model (e.g., Weertman & Weertman, 1964; J. C. Savage & Burford, 1973; Sandwell, 2022), with posterior distributions shown in Figs. S16 and S17.

A key finding from our analysis is that a significant portion of apparent velocity gradients across some faults can be attributed to surface creep, particularly along the Izmit and İsmetpaşa segments of the NAF, where near-fault GNSS station coverage is dense. The inset in the bottom right of Fig. 5d compares strain rates from models that either explicitly incorporate or disregard velocity discontinuities due to surface fault creep, revealing that strain rates exceeding 300 nanostrain/yr may be associated with mismodeled strain rates arising from surface creep along the İsmetpaşa segment. This distinction is important because previous strain rate models in Anatolia did not explicitly account for fault creep, thereby failing to capture the velocity discontinuities observed across faults creeping at the surface—a challenge previously highlighted by Weiss et al. (2020). Differentiating between interseismic fault locking and surface creep contributions to surface strain rates is crucial for seismic hazard assessment, as only strain rates from elastic coupling on faults contribute to the energy budget available for future earthquakes.

Finally, Fig. 6c presents the distribution of mean posterior shallow creep rates along various fault segments in the eastern Mediterranean, compared with prior distributions from previous geodetic studies (e.g., Cakir et al., 2005; Kaneko et al., 2013; Cetin et al., 2014; Cavalié & Jónsson, 2014; Hussain et al., 2016; Hamiel et al., 2016; Yamamoto et al., 2019; Aslan et al., 2019; Bletery et al., 2020; Özarpacı et al., 2021; Hamiel & Piatibratova, 2021; Kurt et al., 2023; Cakir et al., 2023). Our inversion applied moderate constraints on the prior distribution of surface creep rates, allowing the posterior to vary from the prior but ensuring consistency with existing observations where available. The validity of this approach is demonstrated by the minimal and non-systematic residual GNSS velocities across well-instrumented creeping faults, such as the Izmit and İsmetpaşa




Figure 6. (a) Comparison of observed velocities (blue vectors) and mean posterior velocities (red vectors). Profiles 1 and 2 show observed, mean posterior, and screw-dislocation model-predicted velocities (Weertman & Weertman, 1964; J. C. Savage & Burford, 1973) across the İsmetpaşa and Düzce segments of the North Anatolian Fault. Profiles demonstrate accurate modeling of velocity gradients —even across creeping faults— when dense near-fault GNSS velocities are available. (b) Residual velocities (observed minus modeled) colored by their systematic misfit index (red: systematic, blue: random). Histograms show the distribution of east and north velocity residuals. (c) Mean posterior creep rates on select fault segments (red: right-lateral creep rates, blue: left-lateral creep rates), with a profile showing prior and mean posterior creep rate variability along select fault segments as a function of longitude.

segments of the North Anatolian Fault, as shown in Fig. 6b and profile 1 in Fig. 6a. However, surface creep rates along the Palu segment of the East Anatolian Fault and the offshore Marmara segment of the North Anatolian Fault are less well constrained due to limited near-fault GNSS data along these segments.

#### 4.3.2 India-Asia Collision Zone

569

570

571

573

574

575

576

579

580

581

582

583

584

587

588

589

590

591

592

593

595

596

599

600

601

602

603

607

608

610

611

612

613

614

615

616

618

619

620

The ongoing convergence of the Indian and Eurasian plates since the Eocene ( $\sim 50$ Ma) has driven the uplift, deformation, and lateral escape of the Tibetan Plateau (e.g., Molnar & Tapponnier, 1975; Clark & Royden, 2000; Yin & Harrison, 2000; Searle et al., 2011; M. Wang & Shen, 2020; S. Liu et al., 2025). This region has long been a focal point of research aimed at understanding the mechanisms governing present-day horizontal and vertical deformation, as well as its temporal evolution (Thatcher, 2009). Debate over these processes has centered around two end-member models: (1) deformation localized along discrete faults (e.g., Tapponnier et al., 1982; Peltzer & Tapponnier, 1988; Avouac & Tapponnier, 1993; Tapponnier et al., 2001; Replumaz & Tapponnier, 2003; Z.-K. Shen et al., 2005; Meade, 2007; Thatcher, 2007; Y. Li et al., 2025), and (2) deformation occurring as a broadly distributed continuum (e.g., Vilotte et al., 1986; Houseman & England, 1993; Holt et al., 2000; Clark & Royden, 2000; Flesch et al., 2001; England & Molnar, 2005). The first model suggests that most surface deformation is accommodated along a finite number of faults bounding rigid tectonic blocks, whereas the second model posits that strain is diffusely distributed over large regions. An intermediate hypothesis reconciles these two perspectives, proposing that deformation is partitioned between localized faulting and distributed strain within tectonic blocks (e.g., Q. Chen et al., 2004; Loveless & Meade, 2011; Fang et al., 2024).

The first-order characteristics of the GNSS velocity field relative to Eurasia, shown as blue vectors in Fig. 7a, include: (1) a pronounced clockwise rotation of the southeastern borderlands of the Tibetan Plateau around the eastern Himalayan syntaxis, (2) a marked decrease in GNSS velocities across the Himalayas, the Longmen Shan in eastern Tibet, and the Qaidam block in northern Tibet, (3) eastward and westward motion of the Tibetan Plateau's eastern and western flanks, respectively; and (4) a general southeastward motion of lithospheric blocks in eastern Asia relative to the Eurasian Plate, including the South China Block, North China Block, Amurian Plate, and Ordos Block. To assess the quality of our strain rate inversion, we compared observed GNSS velocities with the mean posterior velocities predicted by our model (shown as red vectors in Fig. 7a). The close agreement between the two indicates minimal systematic residuals at regional scales, with roughly 85% of GNSS velocity residuals falling within  $\pm 1 \text{ mm/yr}$ , as illustrated in Fig. 7b. Additionally, a velocity profile across the Xiaojiang Fault compares observed fault-parallel GNSS velocities, model-predicted (mean posterior) velocities from our strain rate inversion, and fault-parallel rates from an antiplane screw dislocation model (e.g., Weertman & Weertman, 1964; J. C. Savage & Burford, 1973), with the posterior distribution shown in Fig. S18. The excellent agreement between model predictions and fault-parallel velocities further reinforces the robustness of our strain rate inversion, accurately capturing velocity gradients across faults (Fig. 7a, panel 1).

While most previous studies interpret GNSS velocities in a Eurasia-fixed reference frame, we find that expressing them in a central Tibet-fixed reference frame (Figs. S5c, S6d) more effectively highlights internal deformation within Tibet and the relative motion of adjacent blocks. This approach, first advocated by Gan et al. (2007), has been largely overlooked in subsequent studies. In this reference frame, the dominant E-W extension across the plateau becomes more apparent. Additionally, it highlights that surface motions around the eastern Himalayan syntaxis cannot be explained by simple rigid block rotation alone. Instead, some internal deformation occurs in the regions bounded by the Jiali and Xianshuihe-Xiaojiang faults, where systematic stretching is clearly evident in the dilatation rate map (Fig. 7c), consistent with previous strain rate studies

(e.g., Allmendinger et al., 2007; Kreemer et al., 2014; M. Wang & Shen, 2020). This deformation pattern was recognized by geologists even before geodetic data became available for crustal deformation studies (e.g., Fig. 1 in Tapponnier et al., 1982) and was attributed to extrusion-related extension. A more detailed analysis of this clockwise rotation was conducted by M. Wang and Shen (2020), who attributed it to a combination of translation and rotation of small blocks within a larger rotating terrane confined between the Jiali and Xianshuihe-Xiaojiang faults. It is important to note, however, that GNSS velocities in this region may still contain postseismic signals from large earthquakes such as the 2001,  $M_W 7.8$  Kunlun and 2008,  $M_W 8.0$  Wenchuan earthquakes, as well as from several smaller earthquakes.

Figure 7c presents the dilatation rates and principal strain rate directions, highlighting key deformation patterns across the region. Significant N-S shortening is observed along the Himalayas, where dilatation rates typically range from -70 to -90 nanostrain/yr, with peaks exceeding -120 nanostrain/yr in the eastern Himalayan syntaxis, consistent with previous studies (e.g., M. Wang & Shen, 2020). Other regions of pronounced shortening include the Hindu Kush (-20 nanostrain/yr), where the maximum shortening direction trends NNE-SSW to NNW-SSE, western Pamir (-60 to -80 nanostrain/yr) and the Pamir Thrust System (-60 to -100 nanostrain/yr), both exhibiting NNW-SSE shortening, Central Tian Shan (-50 to -60 nanostrain/yr) undergoing N-S shortening, and the Qaidam Block in northeastern Tibet, where dilatation rates range from -10 to -20 nanostrain/yr, with principal strain rate directions showing NE-SW shortening. Additionally, NW-SE shortening is observed across the Longmen Shan in eastern Tibet, where typical dilatation rates range from -30 to -40 nanostrain/yr.

In contrast, extension is prominent in the northeastern Pamir, where E-W dilatation rates range from 10 to 40 nanostrain/yr, and across the Tibetan Plateau, where widespread E-W extension results in dilatation rates generally between 10 and 20 nanostrain/yr. The highest dilatation rates within the plateau, around 20 nanostrain/yr, are concentrated in its southern region, where seven N-S oriented rift zones have been documented (Armijo et al., 1986). However, a recent high-resolution strain rate map locally integrating In-SAR and GNSS data revealed that dilatation rates exceed 50 nanostrain/yr along these rift zones (H. Chen et al., 2024), suggesting that strain rates may be higher than inferred from GNSS data alone. This underscores the importance of integrating GNSS and In-SAR observations to refine strain rate estimates and improve the resolution of deformation patterns across this region in future studies. Additionally, a distinct belt of positive dilatation rates is observed between the Jiali and Xianshuihe faults in eastern Tibet, with values typically ranging from 10 to 20 nanostrain/yr, consistent with recent findings by M. Wang and Shen (2020). In this region, principal strain rate directions progressively rotate clockwise around the eastern Himalayan syntaxis.

Our refined strain rate estimates across the India-Asia collision zone and neighboring regions are shown in Fig. 7d, highlighting distinct shear belts. In the westernmost part of our study area, the Chaman Fault exhibits strain rates of 50–60 nanostrain/yr, locally increasing to 70–90 nanostrain/yr in its better-instrumented northern segment. Further north, in the Hindu Kush, strain rates hover around 30 nanostrain/yr. Northward, the Pamir Thrust System shows strain rates between 100 and 150 nanostrain/yr, while the Vakhsh Fault, further west, reaches 130–170 nanostrain/yr. To the west, the Illiac Fault exhibits strain rates of 90–140 nanostrain/yr, decreasing systematically westward. In the Central Tian Shan, northeast of the Pamir Plateau, strain rates typically range from 50 to 60 nanostrain/yr, with local peaks around 100 nanostrain/yr near the Issyk Kul lake.

Moving eastward into the Tibetan Plateau, along the Altyn Tagh Fault, strain rates range from 40 to 60 nanostrain/yr, though GNSS station coverage is sparse. In the southeastern Tarim Basin, where instrumentation is denser, we observe a small cluster of high strain rates (80–150 nanostrain/yr) along the Altyn Tagh Fault. Further south, the Kun-


lun Fault exhibits typical strain rates of 40 nanostrain/yr (see profile D-D', Fig.4), with localized peaks of 60–80 nanostrain/yr, likely influenced by postseismic deformation following the 2001  $M_s$ 8.1 Kunlun earthquake (see Fig. 7d). The Karakorum Fault, in western Tibet, has systematically lower strain rates, around 30 nanostrain/yr. In northeastern Tibet, strain rates along the Haiyuan Fault hover around 60 nanostrain/yr, in agreement with recent results by D.-L. Wu et al. (2024), peaking locally at 80 nanostrain/yr. However, our estimates are lower than the high-resolution InSAR-derived strain rates reported by Ou et al. (2022), which exceed 100 nanostrain/yr, with peaks surpassing 400 nanostrain/yr across a creeping section of the Haiyuan Fault that is not well resolved by our combined GNSS velocity field. Further south, the Ganzi-Yushu Fault displays strain rates of 40–50 nanostrain/yr, while across the Longmen Shan, on the eastern flank of the Tibetan Plateau, strain rates range from 30 to 50 nanostrain/yr.

Across the Himalayas, strain rates consistently exceed 60 nanostrain/yr, with typical values between 90 and 120 nanostrain/yr. The highest strain rates are observed in the eastern Himalayan syntaxis, with strain rates around 140 nanostrain/yr, peaking at 170 nanostrain/yr. Further east, distributed deformation accommodates relative motion between India, Burma, and South China along the southeastern borderlands of the Tibetan Plateau. Here, the Xianshuihe-Xiaojiang Fault System forms a semi-continuous high-strain belt exceeding 100 nanostrain/yr, with localized peaks reaching 180 nanostrain/yr at its northwestern termination, where surface creep has been documented by InSAR studies (e.g., J. Zhang et al., 2018; Y. Li, Nocquet, Shan, & Jian, 2021; Qiao & Zhou, 2021; X. Huang et al., 2023b; Tong & Chen, 2024; Fang et al., 2024).

However, unlike in the Mediterranean, where dense GNSS coverage enabled explicit modeling of along-strike variations in fault creep rates, the irregular and sparse distribution of near-fault GNSS stations along the creeping segments of the Xianshuihe Fault precluded a similar approach (see Fig. S19 in the supplementary material). To the southwest, the Lijiang Fault, oriented NE-SW, forms a narrow shear belt with strain rates of 40–70 nanostrain/yr, peaking above 90 nanostrain/yr. Further south, the Sagaing Fault shows strain rates exceeding 100 nanostrain/yr, in agreement with previous estimates by Lindsey et al. (2023), though slightly lower rates (70–90 nanostrain/yr) are observed in its northern segment due to uneven GNSS station density. Strain rates along the Indo-Burman Ranges similarly range from 50 to 70 nanostrain/yr. Finally, higher posterior uncertainties are observed in regions experiencing greater deformation, particularly along major shear belts such as the Xianshuihe, Xiaojiang, Kunlun, Altyn Tagh, Haiyuan, and Sagaing faults, as well as the Pamir Thrust and Tian Shan, as illustrated in Fig. S14 in the supplementary material.

# 5 Discussion

Building on the extensive GNSS-derived strain rate dataset compiled in this study, we compare our results with other large-scale geophysical datasets indicative of lithospheric and mantle deformation, including stress indicators from the World Stress Map (WSM, Heidbach et al., 2025), seismic strain-rates inferred from the global Centroid Moment Tensor (gCMT) catalog (Dziewonski et al., 1981; Ekström et al., 2012, catalog up to 12/2024), and shear wave splitting (SKS) compilations (Becker et al., 2012, updated as of 02/2025). These comparisons seek to address two fundamental questions: How does the style of strain release from earthquakes compare to the strain accumulation captured by geodetic measurements along the Alpine-Himalayan belt? And to what extent do large-scale mantle deformation patterns, as inferred from seismic anisotropy, correlate with geodetically-derived surface strain rates? To investigate these questions, we conduct three analyses: (1) a comparison between GNSS-derived strain rates and stress orientations from earthquake and borehole data, (2) an assessment of strain rate styles and orientations between geodetic and seismic (Kostrov-summed) strain rate tensors, and (3) an eval-



Top-right histograms display the distribution of East and North GNSS velocity residuals. (c) Dilatation rate (first invariant) and principal strain rate directions (red: extension, blue: shortening). (d) Strain rate. PTS: Pamir Thrust System, VF: Vakhsh Fault, IF: (a) Observed GNSS velocities (blue vectors) compared with mean posterior GNSS velocities (red vectors). Profile 1 illustrates observed, mean postescrew dislocation model-predicted fault-parallel GNSS velocities across the Xiaojiang fault. (b) Residual GNSS velocities (observed-modeled) colored by Illiac Fault, XF: Xianshuihe Fault, XJF: Xiaojiang Fault, IBR: Indo-Burman Ranges systematic, blue: random). their systematic misfit index (red: Figure 7. rior, and

uation of large-scale surface deformation patterns in relation to mantle dynamics inferred from seismic anisotropy.

#### 5.1 Comparison of Strain Rate and Stress Orientations

Although several studies have used the maximum horizontal compressive stress  $(S_{Hmax})$  as a proxy for the maximum shortening rate  $(\dot{\varepsilon}_2)$  direction (e.g., Flesch et al., 2005; Bird, 2009; Z. Shen & Bird, 2022), it has been shown that the two do not always coincide and can exhibit systematic discrepancies in certain regions, such as central Japan, the San Andreas Fault, and the Cascadia subduction zone (e.g., Townend & Zoback, 2006; Schulte-Pelkum et al., 2021; Johnson, 2024).

To investigate this relationship across the Alpide belt, we follow methodologies implemented in previous studies (e.g., Townend & Zoback, 2006; Yang & Hauksson, 2013; Johnson, 2024). In our analysis, we use quality A, B, and C stress data from the World Stress Map (WSM, Heidbach et al., 2025), which are largely derived from earthquake focal mechanisms. As such, they are not entirely independent from strain rates derived from a Kostrov summation, and the relationship between stress and coseismic strain remains subject to ambiguity (e.g., D. P. McKenzie, 1969; Michael, 1987). We identify stress observations within a 50 km radius of each strain rate estimate exceeding 10 nanostrain/yr in the triangular mesh shown in Fig. 3.a. When at least two stress measurements are available, we compute the median stress azimuth and determine its angular difference relative to the maximum shortening direction from our strain rate inversion. Observations are excluded if they do not meet the strain rate threshold, lack sufficient nearby stress data, or if the stress measurements correspond to depths exceeding the local Moho depth from Reguzzoni and Sampietro (2015).

Our results, summarized in Fig. 8, demonstrate a strong large-scale agreement between GNSS-derived maximum shortening rate directions (panel a) and maximum compressive stress orientations (panel b). The azimuthal differences between these parameters are shown in panel c, with the histogram in the bottom-left corner illustrating their distribution. The overall agreement is notable, with a median azimuthal difference of  $-4.1 \pm 15.6$  degrees and a mode of  $-5.0 \pm 15.6$  degrees across the entire Alpide belt, computed using 18 bins (every 10°). Warmer colors (positive values) indicate that the principal shortening rate direction is rotated clockwise relative to  $S_{Hmax}$ , while cooler colors (negative values) represent a counterclockwise offset. Additionally, a violin plot in Fig. S20 of the supplementary material illustrates the distribution of azimuthal differences ( $\dot{\varepsilon}_2 - S_{Hmax}$ ) as a function of the strain rate, highlighting greater variance in lower strain rate regions.

Figure 8 also shows that  $S_{Hmax}$  azimuths exhibit greater spatial variability than  $\dot{\varepsilon}_2$  directions. This difference arises from the nature of the two datasets. The maximum shortening rate azimuths  $(\dot{\varepsilon}_2)$  emphasize broad, coherent deformation patterns, a consequence of the spatial regularization applied in the strain rate inversion. In contrast, the  $S_{Hmax}$  estimates capture local stress conditions and record rotations associated with fault bends, segmentation, and depth variations. Moreover, seismogenic faults are not always optimally oriented relative to the regional stress field (e.g. Townend & Zoback, 2004; Hardebeck & Michael, 2004). Consequently, while the two datasets show strong regional agreement in their median statistics, the  $S_{Hmax}$  azimuths display greater local variability.

# 5.2 Comparison between Geodetic and Seismic Strain Rate Tensors

While comparing stress and principal strain rate directions provides valuable insights into their spatial relationships, a more comprehensive assessment requires analyzing the full strain rate tensor. Here, we evaluate whether strain accumulation and release




Figure 8. (a) GNSS-derived maximum horizontal shortening rate directions ( $\dot{\varepsilon}_2$ ). (b) Maximum horizontal compressive stress ( $S_{Hmax}$ ) azimuths from the World Stress Map (Heidbach et al., 2025), with black lines indicating their orientation. To reduce visual clutter, the WSM dataset has been decimated for display purposes; however, all quality A, B, and C events were included in the analysis. (c) Azimuthal differences between the maximum shortening rate direction and the  $S_{Hmax}$  azimuth ( $\dot{\varepsilon}_2 - S_{Hmax}$ ). The histogram in the bottom-left corner shows the distribution of these azimuthal differences. Warmer colors indicate a clockwise offset of the principal shortening rate direction relative to  $S_{Hmax}$ , while cooler colors indicate an anti-clockwise offset. A mask was applied to show only regions where strain rates exceed 10 nanostrain/yr.

styles are consistent across the Alpide belt by comparing strain rate styles from our GNSS-derived strain rate field with those inferred from Kostrov-summed earthquake moment tensors from the gCMT catalog (Dziewonski et al., 1981; Ekström et al., 2012). As noted, such Kostrov summations are not independent from the WSM stress constraints, the bulk of which are derived from focal mechanisms. However, they are independent from the GNSS-derived strain rates.

ຂດຂ

Following the formulation by Kostrov (1974), the average seismically released strain rate tensor  $\dot{\varepsilon}_{ij}$  within a seismically strained crustal volume V containing N earthquakes with moment tensors  $\mathcal{M}$  over a time period T, given a shear modulus  $\mu$ , is expressed as:

$$\dot{\varepsilon}_{ij} = \frac{1}{2\,\mu\,V\,T} \sum_{n=1}^{N} \mathcal{M}_{ij}^{(n)}.\tag{7}$$

In our analysis, we estimated strain rate tensors by applying a normalized Kostrov summation of moment tensors from the gCMT catalog using an equidistant  $0.5^{\circ} \times 0.5^{\circ}$  grid, considering earthquakes with hypocentral depths shallower than 50 km and  $M_W \geq$  5. We then computed the eigenvalues of both strain rate tensor datasets (Kostrov-summed and GNSS-derived) to obtain the principal strain rates (see details in Appendix A). To ensure spatial consistency, both datasets were linearly interpolated onto a common regular grid, retaining observations where strain rates exceed 10 nanostrain/yr, following the approach used in our previous analysis of stress and principal strain rate orientations.

The strain rate style was computed as a ratio of principal strain rates:  $-\frac{\varepsilon_1+\varepsilon_2}{|\varepsilon_1|+|\varepsilon_2|}$ . This ratio ranges from -1 (indicating a normal faulting style) to +1 (indicating a reverse faulting style), with strike-slip faulting corresponding to values near zero (Johnson, 2024). Figs. 9a and b illustrate the strain rate styles derived from GNSS velocities and Kostrov-summed moment tensors, respectively, while Fig. S21 presents a map of their signed angular differences (GNSS-Kostrov). Strain rate styles derived from Kostrov-summed moment tensors depend on the lower magnitude threshold applied. Due to the Gutenberg-Richter distribution, small earthquakes are more abundant and improve spatial resolution, but their moment tensors are generally less reliable. We tested thresholds ranging from 3.5 (i.e., all events, lowest actual  $M_W$  in the catalog is 4.3) to 6.5 and found that a moment magnitude cutoff of 5 provides a good balance—ensuring sufficient resolution while mitigating the uncertainties associated with smaller earthquake moment tensors. To illustrate this sensitivity, we include strain rate style maps using lower magnitude thresholds of 3.5, 5.5, and 6.5 in the supplementary material (Fig. S22).

Consistent with earlier, more geographically limited comparisons (e.g. Kahle et al., 1998; Holt et al., 2000; Jenny et al., 2004; Faccenna, Becker, Auer, et al., 2014), our results reveal a strong large-scale agreement between strain accumulation and coseismic release styles across the Alpide belt, with a median strain rate style difference of  $0.1\pm0.3$  (Fig. S21). The mode  $\pm$  mean absolute deviation (MAD), computed using 18 bins, matches the median  $\pm$  MAD, reflecting the symmetric distribution of signed strain rate style differences. Small-scale discrepancies primarily arise from the low frequency of instrumental earthquakes along some plate boundaries (e.g., the Himalayas), which biases strain rate styles inferred from Kostrov-summed moment tensors. Additional mismatches occur in subduction settings where intermediate-depth slab earthquakes, such as normal-faulting events in the Caspian Sea linked to slab tearing (e.g., Mumladze et al., 2015) and those within the descending slab of the Makran subduction zone (e.g., Barnhart et al., 2014), influence the inferred strain rate style.

To complement our results, we computed the misfit between geodetic and seismic (Kostrov-summed) strain rate tensors, following a tensor inner product approach used by Frohlich and Davis (1999) and Flesch et al. (2007, eq. 22). To quantify the misfit, we calculated the normalized inner product between the geodetic and seismic strain rate ten-

sors,  $\frac{\dot{\varepsilon}_{geo}:\dot{\varepsilon}_{seis}}{\|\dot{\varepsilon}_{geo}\|\|\dot{\varepsilon}_{seis}\|}$ , which ranges from +1 (perfect match) through 0 (orthogonal) to -1 (opposite orientations). Our results demonstrate large-scale agreement between the orientations of the 2D geodetic and seismic strain rate tensors across the Alpine–Himalayan Belt (Fig. 9c). As noted earlier, opposite geodetic and seismic strain rate orientations occur in the Caspian Sea and inland Makran, where slab-related normal events dominate. Offshore Makran also shows poor agreement, with misfit values close to -1, reflecting both bleeding of our modeled strain rates into unconstrained offshore regions and the influence of normal-faulting earthquakes along the Murray Ridge between the Owen Fracture Zone and the Makran subduction zone. Another area of disagreement lies in the northwestern corner of the North China Block, where geodetic strain rate tensors indicate broadly E-W extension, but most instrumental earthquakes display right-lateral strike-slip motion on NNE-trending faults (Y. G. Zhang et al., 2018).

# 5.3 Large-scale Patterns of GNSS-derived Surface Strain Rates and Mantle Dynamics Inferred from Seismic Anisotropy Compilations

While strain in the crust can be measured directly via geodetic or geological observations, mantle strain must be inferred indirectly from seismic data, either from the earthquake source via moment release or from seismic anisotropy (e.g. Holt, 1995, 2000; Silver, 1996; Silver & Holt, 2002). One of the most widely used methods for inferring mantle strain is through azimuthal seismic anisotropy inferred from shear wave splitting measurements (e.g., Silver & Chan, 1991; Vinnik et al., 1992), as upper mantle anisotropy is largely attributed to crystallographic preferred orientation (CPO) of intrinsically anisotropic olivine crystals (e.g., Christensen & Crosson, 1968; Nicolas & Christensen, 1987; Silver, 1996).

When shear waves pass through an anisotropic medium, they can be split into two orthogonally polarized wave trains that propagate at different speeds. This splitting is quantified by two parameters: the azimuth of the fast-polarized wave ( $\phi$ , "fast axis") and the time delay ( $\delta t$ ) between the fast and slow shear wave components (e.g. Silver, 1996; M. K. Savage, 1999). Core refracted phases such as SK(K)S are often used since this rules out source side anisotropy; while anisotropy may accumulate anywhere along the path, much of the signal is attributed to the uppermost mantle where dislocation creep can form CPOs (e.g. Fischer & Wiens, 1996; Podolefsky et al., 2004; Becker et al., 2008). While the formation and reworking of CPOs is complex, we expect that "A type" fabrics are predominant in the upper mantle, in which case the fast axes should rotate from the maximum extension to the shear orientation for large strains (e.g., S. Zhang & Karato, 1995; Becker et al., 2008, 2014; Bernard et al., 2019). Consequently, SKS fast orientations from seismic anisotropy studies provide valuable insights into lithospheric mantle flow and finite strain (e.g., S. Zhang & Karato, 1995; Silver, 1996).

Early studies comparing SKS fast orientations with surface tectonic features revealed spatial correlations in certain regions. For example, Vinnik et al. (1992) observed that in areas of active plate convergence, SKS fast polarization orientations tend to align with plate boundaries and follow the orientation of associated mountain ranges. Subsequent studies corroborated these findings using strain estimates derived from active faulting and plate-motion models, identifying a general consistency between SKS fast orientations and the directions of maximum crustal horizontal extension in the Tian Shan, Altai, Sayan, and Baikal regions (e.g., Davis et al., 1997). These results suggested coherent mantle and lithospheric deformation and highlighted the potential role of mantle flow in orogenesis (Silver, 1996). In Tibet, McNamara et al. (1994) observed that SKS fast orientations align with fault traces rather than the maximum horizontal elongation directions. Later, with advancements in GNSS instrumentation, Holt (2000) further demonstrated that SKS fast orientations in Tibet systematically align with left-lateral crustal shear planes, supporting the findings of McNamara et al. (1994).




Figure 9. (a) Geodetic strain rate style (red: normal, green: strike-slip, blue: reverse). Moment tensors ( $M_W \geq 5$ , depth < 50 km) are from the gCMT catalog (Dziewonski et al., 1981; Ekström et al., 2012). (b) Seismic strain rate style from Kostrov-summed gCMT tensors with the same criteria. (c) Misfit between geodetic and seismic strain rate tensor orientations, expressed as the dot product of normalized tensors (blue: perfect agreement, green: orthogonal, red: opposite). Results are masked where the strain rate is < 10 nanostrain/yr.

Additionally, several studies identified consistent patterns in rift settings, where SKS fast axes generally parallel the extensional direction, as observed in the Baikal and East African Rift (e.g., Vinnik et al., 1992; Gao et al., 1994; Kendall et al., 2006). Furthermore, SKS fast orientations have been observed to align with the azimuths of major strikeslip faults, such as the Alpine Fault in New Zealand (M. K. Savage et al., 2004) and the strike-slip fault systems of Tibet (e.g., McNamara et al., 1994; Holt, 2000). A similar alignment has also been noted in the upper anisotropic layer required to fit SKS observations along the San Andreas Fault (e.g., Özalaybey & Savage, 1995; Hartog & Schwartz, 2001; M. K. Savage et al., 2004).

However, interpreting shear wave splitting observations remains challenging, particularly in continental regions where complex tectonic histories can lead to significant heterogeneities within the crust and lithosphere. Such heterogeneities can influence both shallow, frozen-in anisotropy and deeper mantle flow (e.g. Silver, 1996; Fouch & Rondenay, 2006; Schulte-Pelkum et al., 2021). Aside from uncertainties surrounding CPO formation, many tectonic scenarios are compatible with multiple interpretations. For instance, observed rift-aligned fast axes may reflect contributions from partial melting, mantle flow, or a combination of both (e.g. Kendall et al., 2006; Faccenna et al., 2013). Moreover, the influence of plate boundaries on rotating fast axes from background trends is not uniform (e.g. Silver & Holt, 2002; Becker et al., 2006; Biryol et al., 2010; Faccenna, Becker, Auer, et al., 2014), with pronounced rotations observed only across faults in few regions. One key issue is that the alignment with crustal features may be indicative of causation, such as a lithospheric origin of anisotropy, or only apparent, such as when the shear in the mantle is coherent with what is observed in the crust (e.g. Silver, 1996; M. K. Savage, 1999).

Our analysis here focuses on documenting the angular relationships between SKS fast orientations  $(\phi)$  and the maximum stretching rate  $(\dot{\varepsilon}_1)$  direction across different sections of the Alpine-Himalayan belt, without attempting to necessarily infer causality or propose specific physical mechanisms behind these correlations. This is achieved with an unprecedented resolution, made possible by our large-scale strain rate estimates, complemented with an up-to-date station-averaged SKS splitting compilation by Becker et al. (2012, updated 02/2025). Fig. 10a provides a large-scale comparison of SKS fast orientations (orange lines) and maximum stretching rate directions (blue lines) across the Alpine-Himalayan belt, with azimuthal difference statistics summarized in Table 3. We report both the median and mode, along with their MAD, to characterize azimuthal differences. However, because the mode more effectively captures dominant azimuthal correlations—particularly in skewed distributions—we adopt the mode  $\pm$  MAD as our primary measure, providing a geophysically meaningful and statistically robust representation of the most frequent azimuthal difference between SKS fast orientations and principal strain rate directions.

#### 5.3.1 Eastern Mediterranean

In the eastern Mediterranean, SKS fast orientations broadly align with GNSS-derived maximum stretching rate directions in the crust  $(\dot{\varepsilon}_1)$ , consistent with previous findings (e.g. Kreemer et al., 2004; Endrun et al., 2011; Merry et al., 2021). Across the region spanning from the Aegean to the Caspian Sea,  $\dot{\varepsilon}_1$  is rotated clockwise relative to SKS, with a mode azimuthal azimuthal difference is  $5.0^{\circ}\pm26.1^{\circ}$  (see histogram 1 in Fig. 10b). In the Aegean, where SKS fast orientations are broadly NE-SW (Hatzfeld et al., 2001; Kreemer et al., 2004; Paul et al., 2014),  $\dot{\varepsilon}_1$  is rotated counterclockwise relative to SKS, with a mode azimuthal difference of  $-15.0^{\circ}\pm6.8^{\circ}$ , as illustrated in histogram 3, Fig. 10c.

Along the North Anatolian Fault (NAF), alignment is particularly strong, with a mode azimuthal difference of  $5.0^{\circ}\pm10.1^{\circ}$ , as shown in histogram 4, Fig. 10c. A systematic along-strike variation is evident: in the eastern NAF,  $\dot{\varepsilon}_1$  is slightly rotated clockwise

**Table 3.** Statistical summary of azimuth differences between SKS fast orientations  $(\phi)$  and maximum principal strain rate  $(\dot{\varepsilon}_1)$  directions in different regions of the Alpine-Himalayan belt.

Region	Data Points	$\mathbf{Median} \pm \mathbf{MAD} \ (°)$	$\mathbf{Mode} \pm \mathbf{MAD} \; (°)$	Map
Eastern Mediterranean	3919	$9.1 \pm 25.5$	$5.0 \pm 26.1$	Fig. 10.b
Tibet and Southeast Asia	3171	$27.7 \pm 22.9$	$25.0 \pm 24.2$	Fig. 10.b
Aegean Sea	146	$-11.5 \pm 6.9$	$-15.0 \pm 6.8$	Fig. 10.c
North Anatolian Fault	694	$6.6 \pm 10.0$	$5.0 \pm 10.1$	Fig. 10.c
Tian Shan	1068	$17.9 \pm 14.4$	$15.0 \pm 14.2$	Fig. 10.c
Altai, Sayan, and Baikal	94	$-5.8 \pm 12.5$	$-5.0 \pm 12.1$	Fig. 10.c

MAD is the median absolute deviation, computed as median  $(|X_i - C(X)|)$ , where C(X) represents either the median or mode of X, depending on which statistics is being presented. The mode was computed using 18 bins (every 10°).

relative to SKS, transitioning westward to a slight counterclockwise rotation, ultimately matching the pattern in the Aegean. In contrast, no clear correlation is detected in central Anatolia; a potential change in alignment of fault proximal azimuthal anisotropy from extensional to maximum shear strain might be expected from fault maturity controlled fabrics (e.g. Biryol et al., 2010; Vauchez et al., 2012; Bonnin et al., 2012) but remains to be further explored. Further south, in the Hellenic and Cyprus back-arc regions, SKS fast orientations and  $\dot{\varepsilon}_1$  are nearly perpendicular, consistent with previous studies (e.g., L. Jolivet et al., 2009, 2013).

#### 5.3.2 Central Asia

924

925

926

927

928

929

930

932

933

934

935

938

939

940

941

942

943

944

945

946

947

948

951

952

953

956

957

958

In the Tian Shan region, SKS fast orientations generally run subparallel to the mountain range, particularly in the western section of the range, with a mode azimuthal difference of  $15.0^{\circ} \pm 14.2^{\circ}$ , in agreement with earlier studies (e.g., Vinnik et al., 1992; Makeyeva et al., 1992; Davis et al., 1997). However, two small clusters of anomalous NNE-SSW SKS fast orientations appear around Issyk Kul and along the Tian Shan–Tarim boundary, resulting in localized azimuthal differences between 50° and 90° between SKS orientations and  $\dot{\varepsilon}_1$ , as shown in Fig. 10.c. These deviations have been attributed to various mechanisms, including small-scale mantle convection (Wolfe & Solomon, 1998), an upper mantle plume (Makeyeva et al., 1992), or shear between the northward-moving lithosphere and the underlying asthenosphere (A. Li & Chen, 2006).

# 5.3.3 Tibet and Southeast Asia

In Tibet and Southeast Asia, SKS fast orientations and  $\dot{\varepsilon}_1$  exhibit a systematic offset of 25.0°  $\pm$  24.2°, with  $\dot{\varepsilon}_1$  consistently rotated clockwise relative to SKS fast orientations across the Tibetan Plateau, as depicted in histogram 2, Fig. 10.b. This result aligns with previous studies (e.g., McNamara et al., 1994; Holt, 2000), which report that SKS fast orientations align with left-lateral crustal shear planes, and are generally subparallel to major strike-slip faults in the region. Additionally, our findings agree with M. K. Savage (1999), who first documented a systematic misalignment of 10°–30° between SKS fast orientations and maximum elongation orientations from plate motion models, with the latter exhibiting a consistent clockwise rotation relative to the SKS fast orientations.

A notable regional variation occurs in Yunnan and Indochina, where SKS fast orientations transition from N-S to E-W. This sharp change has been used to assess the extent of inferred viscous crustal channel flow, suggesting crustal and mantle deformation may be coupled in Southeast Tibet but decoupled farther south in Yunnan and Indochina (e.g., Royden et al., 1997; Clark & Royden, 2000; Flesch et al., 2005; Lev et al., 2006; Kong et al., 2018; Lin & Zhao, 2024). Additionally, this shift has been linked to a pos-




Figure 10. (a) GNSS-derived maximum stretching rate azimuths  $\dot{\varepsilon}_1$  (blue) and SKS fast orientations (orange), compiled by Becker et al. (2012, updated 02/2025). (b,c) Signed azimuthal differences between  $\dot{\varepsilon}_1$  and SKS fast orientations, with histograms showing distributions across the Alpide Belt (top-left) and in selected regions: (1) eastern Mediterranean, (2) Tibet and SE Asia, (3) Aegean Sea, (4) North Anatolian fault, (5) Tian Shan, and (6) Altai-Sayan-Baikal regions. Dashed vertical gray lines mark maximum shear strain rate directions ( $\pm 45^{\circ}$  from  $\dot{\varepsilon}_1$ ,  $\dot{\varepsilon}_2$ ). Warm colors indicate clockwise offsets of  $\dot{\varepsilon}_1$  relative to fast SKS directions, cool colors counterclockwise, and dark purple near-perpendicular orientations. A mask was applied to show only regions where strain rates exceed 10 nanostrain/yr.

sible change in upper mantle deformation style, transitioning from simple shear in Southeast Tibet to pure shear in Yunnan (e.g., C.-Y. Wang et al., 2008).

#### 5.3.4 Altai, Sayan and Baikal Regions

Despite limited observations, we find a general alignment between SKS fast polarization and  $\dot{\varepsilon}_1$  in the Altai, Sayan, and Baikal regions, with a mode azimuthal difference of  $-5.0^{\circ} \pm 12.1^{\circ}$ , as shown in histogram 6 of Fig. 10.c, consistent with earlier findings (e.g., Vinnik et al., 1992; Gao et al., 1994; Davis et al., 1997). Our results thus reinforce and further substantiate previous studies documenting systematic regional-scale correlations between principal strain rate orientations and SKS fast orientations from seismic anisotropy across the Alpide belt. How much of this alignment arises due to lithospheric and how much due to mantle contributions needs to be further explored by tectonic and mantle flow modeling.

## 6 Conclusion

We estimate strain rates along the Alpine-Himalayan belt using a unified, comprehensive, and up-to-date GNSS velocity field constructed with FICORO\_GNSS, a Python toolbox developed for filtering and combining GNSS velocities. Our results show a remarkable agreement between both the styles and orientations of geodetic and seismic (Kostrov-summed) strain rate tensors. Similarly, we observe a strong correspondence between strain rate and stress orientations in the crust. These results suggest overall consistency between the styles of strain accumulation and release across the Alpide belt.

Additionally, we find that mismodeled strain rates due to surface fault creep can be substantial in some regions—for example, exceeding 300 nanostrain/yr along the İsmetpaşa segment of the North Anatolian Fault. To avoid misattributing these localized velocity discontinuities to strain rates from elastic coupling on faults, we explicitly incorporate surface creep rates into our inversion where dense near-fault geodetic data are available. Distinguishing between strain rates associated with elastic coupling on faults and those arising from surface fault creep is essential for seismic hazard assessment, as only the former contributes to the elastic energy buildup to be released in future earthquakes.

Our analyses substantiate previous findings of regional-scale relationships between surface deformation and SKS fast orientations from seismic anisotropy compilations, which are thought to be associated with different mantle deformation regimes. While these correlations do not necessarily imply causation and remain poorly understood, we hypothesize that the observed systematic angular relationships between SKS and principal strain rate directions in regions such as the Aegean, North Anatolia, Tibet, Tian Shan, Altai, Sayan, and Baikal, may reflect deep-seated mantle dynamics, as has been previously suggested. Nevertheless, this hypothesis warrants further investigation, particularly regarding the effects of depth-dependent anisotropy in the lithosphere and its relationship with surface deformation.

# 7 Open Research

All results and figures presented in this study are fully reproducible using the Jupyter notebook and supporting files archived in an open-access repository (Castro-Perdomo, 2025). The Python toolkit FICORO\_GNSS Version 1.3.0, developed for filtering and combining GNSS velocity fields, is distributed under the MIT License and permanently archived in a public repository (Castro-Perdomo, 2024). The MATLAB code BForStrain Version 1.1.0 was used to invert GNSS velocities for strain rates and is publicly available in an open-access repository (Johnson, 2023). Comparisons between GNSS-derived strain rates and  $S_{Hmax}$  orientations were conducted using the World Stress Map (WSM 2025) dataset, available through the GFZ Data Services Portal (Heidbach et al., 2025). Maps were gen-

erated using the Generic Mapping Tools (GMT) Version 6.5.0 (Wessel et al., 2019), PyGMT Version 0.12.0 (Tian et al., 2024), and the scientific color maps of Crameri (2023).

# Acknowledgments

We thank Romain Jolivet and Jorge Jara for providing GNSS velocities for the İsmetpaşa segment of the North Anatolian Fault, and Seda Özarpacı for contributing GNSS velocities across the Izmit segment. We are also grateful to Jeff Freymueller and an anonymous reviewer for their constructive comments, which helped improve the manuscript, and to Michael Hamburger for his thoughtful in-house reviews. This research was supported by the National Science Foundation (NSF) under award EAR-2045292, and in part by King Abdullah University of Science and Technology (KAUST) under award OSR-CRG2020-4335.

# Appendix A The 2D Strain Rate Tensor and its Invariants

Given a set of geodetically-derived horizontal surface velocities v, we can define the strain rate tensor as:

$$\dot{\varepsilon}_{i,j} = \frac{1}{2} \left( \frac{\partial v_i}{\partial x_j} + \frac{\partial v_j}{\partial x_i} \right) = \begin{pmatrix} \frac{\partial v_1}{\partial x_1} & \frac{1}{2} \left( \frac{\partial v_1}{\partial x_2} + \frac{\partial v_2}{\partial x_1} \right) \\ \frac{1}{2} \left( \frac{\partial v_2}{\partial x_1} + \frac{\partial v_1}{\partial x_2} \right) & \frac{\partial v_2}{\partial x_2} \end{pmatrix}, \tag{A1}$$

where i and j represent the components of coordinates x and surface velocities v. Here, we consider surface horizontal strain rates, then i, j = 1, 2, where the eigenvalues of the strain rate tensor represent the principal strain rates ( $\dot{\varepsilon}_1$  and  $\dot{\varepsilon}_2$ ):

$$\dot{\varepsilon}_1, \dot{\varepsilon}_2 = \frac{\dot{\varepsilon}_{11} + \dot{\varepsilon}_{22}}{2} \pm \sqrt{\left(\frac{\dot{\varepsilon}_{11} - \dot{\varepsilon}_{22}}{2}\right)^2 + \dot{\varepsilon}_{12}^2},\tag{A2}$$

and their corresponding eigenvectors are the principal strain rate orientations. In continuum mechanics, we often use the tensor's coordinate system independent quantities (i.e., the invariants) for quantifying geodetically-derived surface strain rates. These invariants appear as coefficients of the characteristic polynomial of the tensor, which for a 2D horizontal strain rate tensor is given by  $\lambda^2 - I_1(\dot{\varepsilon}) \lambda + I_2(\dot{\varepsilon}) = 0$ , where  $I_1$  and  $I_2$  are the first and second invariants, respectively.

The first invariant is the trace of the strain rate tensor, representing the areal strain rate (dilatation rate,  $\dot{\Delta}$ ), which is equivalent to the divergence of the velocity field:

$$I_1^{2D}(\dot{\varepsilon}) = \dot{\Delta} = \nabla \cdot v = \operatorname{tr}(\dot{\varepsilon}) = \dot{\varepsilon}_{11} + \dot{\varepsilon}_{22} = \dot{\varepsilon}_1 + \dot{\varepsilon}_2. \tag{A3}$$

A related quantity is the mean (normal) horizontal strain rate  $(\dot{\varepsilon}_m^{2D})$ , defined as:

$$\dot{\varepsilon}_m^{2D} = \frac{1}{2}I_1^{2D} = \frac{1}{2}(\dot{\varepsilon}_1 + \dot{\varepsilon}_2) \tag{A4}$$

The second invariant can be expressed in several equivalent forms. The formal definition can be written as:

$$I_2(\dot{\varepsilon}) = \frac{1}{2} \left[ \operatorname{tr}(\dot{\varepsilon})^2 - \operatorname{tr}(\dot{\varepsilon}^2) \right]. \tag{A5}$$

In the special case of a 2D tensor, this reduces directly to the determinant:

1035

1045

1046

1047

1048

$$I_2^{2D}(\dot{\varepsilon}) = \det(\dot{\varepsilon}) = \dot{\varepsilon}_{11}\dot{\varepsilon}_{22} - \dot{\varepsilon}_{12}^2.$$
 (A6)

Equivalently, because the determinant of a tensor equals the product of its eigenvalues, the second invariant of the 2D strain rate tensor is also:

$$I_2^{2D}(\dot{\varepsilon}) = \dot{\varepsilon}_1 \,\dot{\varepsilon}_2. \tag{A7}$$

The definition of the second invariant of the 2D horizontal strain rate tensor has been applied inconsistently in the literature. A commonly used expression is:

$$\|\dot{\varepsilon}\|_F^{2D} = \sqrt{\dot{\varepsilon}_{11}^2 + \dot{\varepsilon}_{22}^2 + 2\dot{\varepsilon}_{12}^2},$$

$$\|\dot{\varepsilon}\|_F^{2D} = \sqrt{\dot{\varepsilon}_1^2 + \dot{\varepsilon}_2^2},$$
(A8)

which corresponds to the Frobenius (Euclidean) norm of the 2D horizontal strain rate tensor rather than its formal second invariant  $(I_2, \text{Eq. A5})$ . The Frobenius norm is often preferred because it is always non-negative, whereas the formal second invariant  $(I_2)$  can be negative if  $\dot{\varepsilon}_{11}\dot{\varepsilon}_{22}-\dot{\varepsilon}_{12}^2<0$ . To facilitate direct comparison with previous studies, we adopted the Frobenius norm in this work.

Another common measure is the maximum shear strain rate  $(\dot{\gamma}_{max})$ , which relates to the second invariant of the deviatoric strain rate tensor  $(J_2)$ , and in 2D is given by:

$$|\dot{\gamma}_{max}^{2D}| = \sqrt{J_2}, \dot{\gamma}_{max}^{2D} = \frac{1}{2}\sqrt{(\dot{\varepsilon}_{11} - \dot{\varepsilon}_{22})^2 + 4\dot{\varepsilon}_{12}} = \frac{1}{2}(\dot{\varepsilon}_1 - \dot{\varepsilon}_2).$$
(A9)

Combining equations A2 and A9, we find that the principal extension and shortening rates  $\dot{\varepsilon_1}, \dot{\varepsilon_2}$  can be rewritten in terms of  $\dot{\Delta}$  and  $\dot{\gamma}_{max}$ 

$$\dot{\varepsilon}_{1} = \frac{\dot{\Delta} + \dot{\gamma}_{max}}{2},$$

$$\dot{\varepsilon}_{2} = \frac{\dot{\Delta} - \dot{\gamma}_{max}}{2}.$$

A relationship exists between  $J_2$ ,  $I_1 = \operatorname{tr}(\dot{\varepsilon})$ , and  $\|\dot{\varepsilon}\|_F^{2D}$ , which can be written as:

$$\|\dot{\varepsilon}\|_F^{2D} = \sqrt{2J_2 + \frac{1}{2}(I_1)^2}.$$
 (A10)

From eq. A10, it is clear that for a 2D deviatoric strain rate tensor  $\dot{\varepsilon}'$ , the Frobenius norm is just a scaled form of the deviatoric second invariant, or equivalently, a scaled form of the maximum shear strain rate:

$$\|\dot{\varepsilon}'\|_F^{2D} = \sqrt{2J_2} = \sqrt{2} \cdot \dot{\gamma}_{max}^{2D}.$$
 (A11)

The azimuth of the maximum principal strain rate  $(\dot{\varepsilon}_{max})$  can be estimated as follows:

$$Az_{\dot{\varepsilon}_{\max}} = 90^{\circ} - \frac{\arctan\left(\frac{\dot{\varepsilon}_{12}}{\frac{\dot{\varepsilon}_{11} - \dot{\varepsilon}_{22}}{2}}\right)}{2}.$$
 (A12)

We can further estimate the azimuth of the maximum shortening rate  $(\dot{\varepsilon}_2)$  by evaluating the sign of the maximum principal strain rate  $(\dot{\varepsilon}_{\max})$ . If  $\dot{\varepsilon}_{\max} > 0$  (extension), the maximum shortening rate occurs 90° away from the azimuth of  $\dot{\varepsilon}_{\max}$ . On the other hand, if  $\dot{\varepsilon}_{\max} < 0$  (compression), the azimuth of  $\dot{\varepsilon}_{\max}$  directly represents the maximum shortening direction.

# References

- Abdrakhmatov, K., Arrowsmith, R., Elliott, J., Grutzner, C., Mukambayev, A., Rizza, M., . . . Wilkinson, R. (2024). Urgent need for greater earthquake resilience in continental Asia. *Nature Geoscience*, 17(9), 818–819. doi: https://doi.org/10.1038/s41561-024-01531-0
- Alchalbi, A., Daoud, M., Gomez, F., McClusky, S., Reilinger, R., Romeyeh, M. A., ... Barazangi, M. (2010, January). Crustal deformation in northwestern Arabia from GPS measurements in Syria: Slow slip rate along the northern Dead Sea Fault. Geophysical Journal International, 180(1), 125–135. Retrieved 2019-06-21, from https://academic.oup.com/gji/article-lookup/doi/10.1111/j.1365-246X.2009.04431.x doi: 10.1111/j.1365-246X.2009.04431.x
- Allmendinger, R. W., Reilinger, R., & Loveless, J. (2007). Strain and rotation rate from GPS in Tibet, Anatolia, and the Altiplano. *Tectonics*, 26(3). doi: https://doi.org/10.1029/2006TC002030
- Altamimi, Z., Métivier, L., Rebischung, P., Rouby, H., & Collilieux, X. (2017, June). ITRF2014 plate motion model. Geophysical Journal International, 209(3), 1906-1912. Retrieved 2018-07-17, from https://academic.oup.com/gji/article-lookup/doi/10.1093/gji/ggx136 doi: 10.1093/gji/ggx136
- Altamimi, Z., Sillard, P., & Boucher, C. (2002). ITRF2000: A new release of the International Terrestrial Reference Frame for earth science applications. *Journal of Geophysical Research: Solid Earth*, 107(B10), ETG 2-1-ETG 2-19. doi: https://doi.org/10.1029/2001JB000561
- Argand, E. (1922). La tectonique de l'Asie. Conférence faite á Bruxelles, le 10 août 1922. In Congrès géologique international (XIIIe session)-Belgique 1922 (pp. 171–372).
- Armijo, R., Tapponnier, P., Mercier, J. L., & Han, T.-L. (1986). Quaternary extension in southern Tibet: Field observations and tectonic implications. *Journal of Geophysical Research: Solid Earth*, 91 (B14), 13803-13872. doi: https://doi.org/10.1029/JB091iB14p13803
- Aslan, G., Lasserre, C., Cakir, Z., Ergintav, S., Özarpaci, S., Dogan, U., ... Renard, F. (2019). Shallow Creep Along the 1999 Izmit Earthquake Rupture (Turkey) From GPS and High Temporal Resolution Interferometric Synthetic Aperture Radar Data (2011–2017). *Journal of Geophysical Research: Solid Earth*, 124(2), 2218-2236. doi: https://doi.org/10.1029/2018JB017022

Avouac, J.-P., & Tapponnier, P. (1993). Kinematic model of active deformation in central Asia. *Geophysical Research Letters*, 20 (10), 895-898. doi: https://doi.org/10.1029/93GL00128

- Bahrouni, N., Masson, F., Meghraoui, M., Saleh, M., Maamri, R., Dhaha, F., & Arfaoui, M. (2020). Active tectonics and GPS data analysis of the Maghrebian thrust belt and Africa-Eurasia plate convergence in Tunisia. *Tectonophysics*, 785, 228440. doi: https://doi.org/10.1016/j.tecto.2020.228440
- Bai, D., Unsworth, M. J., Meju, M. A., Ma, X., Teng, J., Kong, X., . . . others (2010). Crustal deformation of the eastern Tibetan plateau revealed by magnetotelluric imaging. *Nature geoscience*, 3(5), 358–362. doi: https://doi.org/10.1038/ngeo830
- Barbot, S., & Weiss, J. (2021). Connecting subduction, extension and shear localization across the Aegean Sea and Anatolia. *Geophysical Journal International*, 226(1), 422–445. doi: https://doi.org/10.1093/gji/ggab078
- Barnhart, W. D. (2017). Fault creep rates of the Chaman fault (Afghanistan and Pakistan) inferred from InSAR. *Journal of Geophysical Research: Solid Earth*, 122(1), 372-386. doi: https://doi.org/10.1002/2016JB013656
- Barnhart, W. D., Hayes, G. P., Samsonov, S. V., Fielding, E. J., & Seidman, L. E. (2014). Breaking the oceanic lithosphere of a subducting slab: The 2013 Khash, Iran earthquake. Geophysical Research Letters, 41(1), 32-36. doi: https://doi.org/10.1002/2013GL058096
- Beavan, J., & Haines, J. (2001). Contemporary horizontal velocity and strain rate fields of the Pacific-Australian plate boundary zone through New Zealand.

  Journal of Geophysical Research: Solid Earth, 106(B1), 741–770.
- Becker, T. W., Conrad, C. P., Schaeffer, A. J., & Lebedev, S. (2014). Origin of azimuthal seismic anisotropy in oceanic plates and mantle. *Earth Planet. Sci. Lett.*, 401, 236–250.
- Becker, T. W., & Faccenna, C. (2011). Mantle conveyor beneath the Tethyan collisional belt. Earth and Planetary Science Letters, 310(3), 453-461. doi: https://doi.org/10.1016/j.epsl.2011.08.021
- Becker, T. W., Kustowski, B., & Ekström, G. (2008). Radial seismic anisotropy as a constraint for upper mantle rheology. *Earth Planet. Sci. Lett.*, 267, 213–237.
- Becker, T. W., Lebedev, S., & Long, M. D. (2012). On the relationship between azimuthal anisotropy from shear wave splitting and surface wave tomography. *Journal of Geophysical Research: Solid Earth*, 117(B1). doi: https://doi.org/10.1029/2011JB008705
- Becker, T. W., Schulte-Pelkum, V., Blackman, D. K., Kellogg, J. B., & O'Connell, R. J. (2006). Mantle flow under the western United States from shear wave splitting. Earth and Planetary Science Letters, 247(3), 235-251. doi: https://doi.org/10.1016/j.epsl.2006.05.010
- Bernard, R., Behr, W. M., Becker, T. W., & Young, D. (2019). Relationships between olivine CPO and deformation parameters in naturally deformed rocks and implications for mantle seismic anisotropy. *Geochem.*, *Geophys.*, *Geosys.*, 20, 3469–3494.
- Bilham, R. (2006). Dangerous Tectonics, Fragile Buildings, and Tough Decisions. *Science*, 311(5769), 1873-1875. doi: 10.1126/science.1125176
- Bilham, R., Ozener, H., Mencin, D., Dogru, A., Ergintav, S., Cakir, Z., ... Mattioli, G. (2016). Surface creep on the North Anatolian Fault at Ismetpasa, Turkey, 1944–2016. Journal of Geophysical Research: Solid Earth, 121(10), 7409-7431. doi: https://doi.org/10.1002/2016JB013394
- Billi, A., Cuffaro, M., Orecchio, B., Palano, M., Presti, D., & Totaro, C. (2023).

  Retracing the Africa–Eurasia nascent convergent boundary in the Western

  Mediterranean based on earthquake and GNSS data. Earth and Planetary

  Science Letters, 601, 117906.
  - Bird, P. (2009). Long-term fault slip rates, distributed deformation rates, and

forecast of seismicity in the western United States from joint fitting of community geologic, geodetic, and stress direction data sets. *Journal of Geophysical Research: Solid Earth*, 114 (B11). doi: 10.1029/2009JB006317

- Biryol, C. B., Zandt, G., Beck, S. L., Ozacar, A. A., Adiyaman, H. E., & Gans, C. R. (2010). Shear wave splitting along a nascent plate boundary: the North Anatolian Fault Zone. *Geophys. J. Int.*, 181, 1201–1213.
- Bletery, Q., Cavalié, O., Nocquet, J.-M., & Ragon, T. (2020). Distribution of Interseismic Coupling Along the North and East Anatolian Faults Inferred From InSAR and GPS Data. *Geophysical Research Letters*, 47(16), e2020GL087775. doi: https://doi.org/10.1029/2020GL087775
- Bonnin, M., Tommasi, A., Hassani, R., Chevrot, S., Wookey, J., & Barruol, G. (2012). Numerical modelling of the upper-mantle anisotropy beneath a migrating strike-slip plate boundary: the San Andreas Fault system. *Geophys. J. Int.*, 191, 436–458.
- Bougrine, A., Yelles-Chaouche, A. K., & Calais, E. (2019). Active deformation in Algeria from continuous GPS measurements. *Geophysical Journal International*, 217(1), 572–588.
- Briole, P., Ganas, A., Elias, P., & Dimitrov, D. (2021). The GPS velocity field of the Aegean. New observations, contribution of the earthquakes, crustal blocks model. *Geophysical Journal International*, 226(1), 468–492.
- Brockmann, E., Lutz, S., Zurutuza, J., Caporali, A., Lidberg, M., Völksen, C., ... others (2019). Towards a Dense Velocity Field in Europe as a basis for Maintaining the European Reference Frame. 27th IUGG Assembly, Montreal.
- Brückl, E., Bleibinhaus, F., Gosar, A., Grad, M., Guterch, A., Hrubcová, P., . . . Thybo, H. (2007). Crustal structure due to collisional and escape tectonics in the Eastern Alps region based on profiles Alp01 and Alp02 from the ALP 2002 seismic experiment. Journal of Geophysical Research: Solid Earth, 112 (B6). doi: https://doi.org/10.1029/2006JB004687
- Cai, J., & Grafarend, E. W. (2007). Statistical analysis of geodetic deformation (strain rate) derived from the space geodetic measurements of BIFROST Project in Fennoscandia. *Journal of Geodynamics*, 43(2), 214–238.
- Cakir, Z., Akoglu, A. M., Belabbes, S., Ergintav, S., & Meghraoui, M. (2005).
  Creeping along the Ismetpasa section of the North Anatolian fault (Western Turkey): Rate and extent from InSAR. Earth and Planetary Science Letters, 238(1), 225-234. doi: https://doi.org/10.1016/j.epsl.2005.06.044
- Cakir, Z., Doğan, U., Akoğlu, A. M., Ergintav, S., Özarpacı, S., Özdemir, A., ... Bilham, R. (2023). Arrest of the Mw 6.8 January 24, 2020 Elaziğ (Turkey) earthquake by shallow fault creep. Earth and Planetary Science Letters, 608, 118085. doi: https://doi.org/10.1016/j.epsl.2023.118085
- Carey, S. W. (1955). The orocline concept in geotectonics-Part I. Papers and Proceedings of the Royal Society of Tasmania.
- Carminati, E., Doglioni, C., & Scrocca, D. (2003). Apennines subduction-related subsidence of Venice (Italy). Geophysical Research Letters, 30(13). doi: https://doi.org/10.1029/2003GL017001
- Castro-Perdomo, N. (2024). FICORO\_GNSS: An open-source Python software package for filtering, combining and rotating GNSS velocity fields (Version 1.3.0) [Software]. Zenodo. doi: https://doi.org/10.5281/zenodo.13921188
- Castro-Perdomo, N. (2025). Supplementary material for "Strain Rates along the Alpine-Himalayan Belt from a Comprehensive GNSS Velocity Field" [Dataset]. Zenodo. doi: https://doi.org/10.5281/zenodo.15186333
- Castro-Perdomo, N., Viltres, R., Masson, F., Klinger, Y., Liu, S., Dhahry, M., ... others (2022). Interseismic deformation in the Gulf of Aqaba from GPS measurements. *Geophysical Journal International*, 228(1), 477–492.
- Cavalié, O., & Jónsson, S. (2014). Block-like plate movements in eastern Anatolia observed by InSAR. Geophysical Research Letters, 41(1), 26-31. doi: https://

doi.org/10.1002/2013GL058170

- Cetin, E., Cakir, Z., Meghraoui, M., Ergintav, S., & Akoglu, A. M. (2014). Extent and distribution of aseismic slip on the Ismetpaşa segment of the North Anatolian Fault (Turkey) from Persistent Scatterer InSAR. Geochemistry, Geophysics, Geosystems, 15(7), 2883-2894. doi: https://doi.org/10.1002/2014GC005307
- Chen, H., Qu, C., Zhao, D., Shan, X., Li, C., & Dal Zilio, L. (2024). Large-Scale Extensional Strain in Southern Tibet From Sentinel-1 InSAR and GNSS Data. *Geophysical Research Letters*, 51(19), e2024GL110512. doi: https://doi.org/10.1029/2024GL110512
- Chen, Q., Freymueller, J. T., Wang, Q., Yang, Z., Xu, C., & Liu, J. (2004, 1). A deforming block model for the present-day tectonics of Tibet. Journal of Geophysical Research: Solid Earth, 109. doi: 10.1029/2002JB002151
- Chousianitis, K., Sboras, S., Mouslopoulou, V., Chouliaras, G., & Hristopulos, D. T. (2024). The Upper Crustal Deformation Field of Greece Inferred From GPS Data and Its Correlation With Earthquake Occurrence. *Journal of Geophysical Research: Solid Earth*, 129(4), e2023JB028004. doi: https://doi.org/10.1029/2023JB028004
- Christensen, N., & Crosson, R. (1968). Seismic anisotropy in the upper mantle. Tectonophysics, 6(2), 93-107. doi: https://doi.org/10.1016/0040-1951(68)90013-9
- Clark, M. K., & Royden, L. H. (2000). Topographic ooze: Building the eastern margin of Tibet by lower crustal flow. *Geology*, 28, 703–706.
- Crameri, F. (2023). Scientific colour maps (Version 8.0.1) [Software]. Zenodo. Retrieved from https://doi.org/10.5281/zenodo.8409685 doi: 10.5281/zenodo.8409685
- Danciu, L., Giardini, D., Weatherill, G., Basili, R., Nandan, S., Rovida, A., ... Wiemer, S. (2024). The 2020 European Seismic Hazard Model: Overview and Results. *EGUsphere*, 2024, 1–36. doi: 10.5194/egusphere-2023-3062
- Davis, P., England, P., & Houseman, G. (1997). Comparison of shear wave splitting and finite strain from the India-Asia collision zone. *Journal of Geophysical Research: Solid Earth*, 102(B12), 27511-27522. doi: https://doi.org/10.1029/97JB02378
- Dercourt, J. e. a., Zonenshain, L., Ricou, L.-E., Kazmin, V., Le Pichon, X., Knipper, A., ... others (1986). Geological evolution of the Tethys belt from the Atlantic to the Pamirs since the Lias. *Tectonophysics*, 123(1-4), 241–315.
- Devoti, R., d'Agostino, N., Serpelloni, E., Pietrantonio, G., Riguzzi, F., Avallone, A., . . . others (2017). A combined velocity field of the Mediterranean region. Annals of Geophysics, 60(2). doi: 10.4401/ag-7059
- DeVries, P. M., Krastev, P. G., Dolan, J. F., & Meade, B. J. (2017). Viscoelastic block models of the North Anatolian fault: A unified earthquake cycle representation of pre-and postseismic geodetic observations. Bulletin of the Seismological Society of America, 107(1), 403–417.
- Dziewonski, A. M., Chou, T.-A., & Woodhouse, J. H. (1981). Determination of earthquake source parameters from waveform data for studies of global and regional seismicity. *Journal of Geophysical Research: Solid Earth*, 86 (B4), 2825–2852.
- Ekström, G., Nettles, M., & Dziewoński, A. (2012). The global CMT project 2004–2010: Centroid-moment tensors for 13,017 earthquakes. *Physics of the Earth and Planetary Interiors*, 200, 1–9.
- El-Fiky, G. S., & Kato, T. (1998). Continuous distribution of the horizontal strain in the Tohoku district, Japan, predicted by least-squares collocation. *Journal of Geodynamics*, 27(2), 213–236.
- Endrun, B., Lebedev, S., Meier, T., Tirel, C., & Friederich, W. (2011). Complex layered deformation within the Aegean crust and mantle revealed by seismic

anisotropy. Nature Geoscience, 4(3), 203-207. doi: https://doi.org/10.1038/ngeo1065

1261

1262

1263

1264

1265

1268

1269

1270

1271

1273

1274

1275

1276

1277

1280

1281

1283

1286

1287

1289

1292

1293

1294

1295

1299

1300

1301

1302

1304

1305

1307

1311

- England, P., & Houseman, G. (1986). Finite strain calculations of continental deformation: 2. Comparison with the India-Asia Collision Zone.

  Journal of Geophysical Research: Solid Earth, 91 (B3), 3664-3676. doi: https://doi.org/10.1029/JB091iB03p03664
- England, P., Houseman, G., & Nocquet, J.-M. (2016). Constraints from GPS measurements on the dynamics of deformation in Anatolia and the Aegean. *Journal of Geophysical Research: Solid Earth*, 121 (12), 8888–8916.
- England, P., & Molnar, P. (2005). Late Quaternary to decadal velocity fields in Asia. *Journal of Geophysical Research: Solid Earth*, 110 (B12). doi: https://doi.org/10.1029/2004JB003541
- Engwirda, D. (2009). Mesh2d-automatic mesh generation. available online on MatlabCentral: http://www.mathworks.com/matlabcentral/fileexchange/25555-mesh2d-automatic-mesh-generation.
- Ergintav, S., Floyd, M., Paradissis, D., Karabulut, H., Vernant, P., Masson, F., ... others (2023). New geodetic constraints on the role of faults and blocks vs. distribute strain in the Nubia-Arabia-Eurasia zone of active plate interactions. *Turkish Journal of Earth Sciences*, 32(3), 248–261.
- Ergintav, S., McClusky, S., Hearn, E., Reilinger, R., Cakmak, R., Herring, T., ... Tari, E. (2009). Seven years of postseismic deformation following the 1999, M = 7.4 and M = 7.2, Izmit-Düzce, Turkey earthquake sequence. Journal of Geophysical Research: Solid Earth, 114(B7). doi: https://doi.org/10.1029/2008JB006021
- Ergintav, S., Vernant, P., Tan, O., Karabulut, H., Özarpacı, S., Floyd, M., ... Farımaz, I. (2024). Unexpected far-field deformation of the 2023 Kahramanmaraş earthquakes revealed by space geodesy. *Science*, 386 (6719), 328-335. doi: 10.1126/science.ado4220
- Faccenna, C., Becker, T. W., Auer, L., Billi, A., Boschi, L., Brun, J. P., . . . Serpelloni, E. (2014). Mantle dynamics in the Mediterranean. *Reviews of Geophysics*, 52(3), 283-332. doi: https://doi.org/10.1002/2013RG000444
- Faccenna, C., Becker, T. W., Jolivet, L., & Keskin, M. (2013). Mantle convection in the Middle East: Reconciling Afar upwelling, Arabia indentation and Aegean trench rollback. *Earth Planet. Sci. Lett.*, 375, 254–269.
- Faccenna, C., Becker, T. W., Miller, M. S., Serpelloni, E., & Willett, S. D. (2014). Isostasy, dynamic topography, and the elevation of the Apennines of Italy. Earth and Planetary Science Letters, 407, 163-174. doi: https://doi.org/10.1016/j.epsl.2014.09.027
- Faccenna, C., Bellier, O., Martinod, J., Piromallo, C., & Regard, V. (2006). Slab detachment beneath eastern Anatolia: A possible cause for the formation of the North Anatolian fault. *Earth Planet. Sci. Lett.*, 242, 85–97.
- Faccenna, C., Molin, P., Orecchio, B., Olivetti, V., Bellier, O., Funiciello, F., . . . Billi, A. (2011). Topography of the Calabria subduction zone (southern Italy): Clues for the origin of Mt. Etna. *Tectonics*, 30(1). doi: https://doi.org/10.1029/2010TC002694
- Fang, J., Wright, T. J., Johnson, K. M., Ou, Q., Styron, R., Craig, T. J., ...

  Zheng, G. (2024). Strain Partitioning in the Southeastern Tibetan Plateau From Kinematic Modeling of High-Resolution Sentinel-1 InSAR and GNSS. Geophysical Research Letters, 51(19), e2024GL111199. doi: https://doi.org/10.1029/2024GL111199
- Fattahi, H., & Amelung, F. (2016). InSAR observations of strain accumulation and fault creep along the Chaman Fault system, Pakistan and Afghanistan.

  Geophysical Research Letters, 43(16), 8399-8406. doi: https://doi.org/10.1002/2016GL070121
  - Fischer, K. M., & Wiens, D. A. (1996). The depth distribution of mantle anisotropy

beneath the Tonga subduction zone. Earth Planet. Sci. Lett., 142, 253–260.

- Flerit, F., Armijo, R., King, G., & Meyer, B. (2004). The mechanical interaction between the propagating North Anatolian Fault and the back-arc extension in the Aegean. Earth and Planetary Science Letters, 224 (3-4), 347–362. doi: https://doi.org/10.1016/j.epsl.2004.05.028
- Flesch, L. M., Haines, A. J., & Holt, W. E. (2001). Dynamics of the India-Eurasia collision zone. *Journal of Geophysical Research: Solid Earth*, 106 (B8), 16435-16460. doi: https://doi.org/10.1029/2001JB000208
- Flesch, L. M., Holt, W. E., Haines, A. J., Wen, L., & Shen-Tu, B. (2007, 06). The dynamics of western North America: stress magnitudes and the relative role of gravitational potential energy, plate interaction at the boundary and basal tractions. *Geophysical Journal International*, 169(3), 866-896. doi: 10.1111/j.1365-246X.2007.03274.x
- Flesch, L. M., Holt, W. E., Silver, P. G., Stephenson, M., Wang, C.-Y., & Chan, W. W. (2005). Constraining the extent of crust—mantle coupling in central Asia using GPS, geologic, and shear wave splitting data. Earth and Planetary Science Letters, 238(1), 248-268. doi: https://doi.org/10.1016/j.epsl.2005.06.023
- Floyd, M., King, R., Paradissis, D., Karabulut, H., Ergintav, S., Raptakis, K., & Reilinger, R. (2023). Variations in coupling and deformation along the Hellenic subduction zone. *Turkish Journal of Earth Sciences*, 32(3), 262–274.
- Fouch, M. J., & Rondenay, S. (2006). Seismic anisotropy beneath stable continental interiors. *Phys. Earth Planet. Inter.*, 158, 292–320.
- Fox, M., Herman, F., Kissling, E., & Willett, S. D. (2015, 05). Rapid exhumation in the Western Alps driven by slab detachment and glacial erosion. *Geology*, 43(5), 379-382. doi: 10.1130/G36411.1
- Frohlich, C., & Davis, S. D. (1999). How well constrained are well-constrained T, B, and P axes in moment tensor catalogs? *Journal of Geophysical Research: Solid Earth*, 104(B3), 4901-4910. doi: https://doi.org/10.1029/1998JB900071
- Fukuda, J., & Johnson, K. M. (2021). Bayesian Inversion for a Stress-Driven Model of Afterslip and Viscoelastic Relaxation: Method and Application to Postseismic Deformation Following the 2011 M 9.0 Tohoku-Oki Earthquake.

  \*\*Journal of Geophysical Research: Solid Earth, 126(5), e2020JB021620. doi: https://doi.org/10.1029/2020JB021620
- Gan, W., Zhang, P., Shen, Z.-K., Niu, Z., Wang, M., Wan, Y., . . . Cheng, J. (2007). Present-day crustal motion within the Tibetan Plateau inferred from GPS measurements. *Journal of Geophysical Research: Solid Earth*, 112(B8). doi: https://doi.org/10.1029/2005JB004120
- Gao, S., Davis, P. M., Liu, H., Slack, P. D., Zorin, Y. A., Mordvinova, V. V., ... Meyer, R. P. (1994). Seismic anisotropy and mantle flow beneath the Baikal rift zone. *Nature*, 371 (6493), 149–151. doi: https://doi.org/10.1038/371149a0
- Gomez, F., Cochran, W. J., Yassminh, R., Jaafar, R., Reilinger, R., Floyd, M., ... Barazangi, M. (2020). Fragmentation of the Sinai Plate indicated by spatial variation in present-day slip rate along the Dead Sea Fault System. *Geophysical Journal International*, 221(3), 1913–1940.
- Gomez, F., Khawlie, M., Tabet, C., Nasser Darkal, A., Khair, K., & Barazangi, M. (2006). Late Cenozoic uplift along the northern Dead Sea transform in Lebanon and Syria. *Earth and Planetary Science Letters*, 241(3), 913-931. doi: https://doi.org/10.1016/j.epsl.2005.10.029
- Graham, S. E., Loveless, J. P., & Meade, B. J. (2018). Global plate motions and earthquake cycle effects. *Geochemistry, Geophysics, Geosystems*, 19(7), 2032–2048.
- Güvercin, S. E., Karabulut, H., Konca, A. Ö., Doğan, U., & Ergintav, S. (2022, 02).

  Active seismotectonics of the East Anatolian Fault. Geophysical Journal International, 230(1), 50-69. doi: 10.1093/gji/ggac045

Haines, A., & Holt, W. (1993). A procedure for obtaining the complete horizontal motions within zones of distributed deformation from the inversion of strain rate data. *Journal of Geophysical Research: Solid Earth*, 98(B7), 12057–12082.

1372

1373

1375

1378

1379

1381

1382

1384

1385

1386

1387

1390

1391

1392

1393

1394

1397

1398

1399

1400

1403

1404

1405

1406

1409

1410

1411

1412

1413

1415

1416

- Haines, A. J., Dimitrova, L. L., Wallace, L. M., & Williams, C. A. (2015). Enhanced surface imaging of crustal deformation: Obtaining tectonic force fields using GPS data. Springer.
  - Hamiel, Y., & Piatibratova, O. (2021). Spatial variations of slip and creep rates along the southern and central Dead Sea Fault and the Carmel–Gilboa Fault System. *Journal of Geophysical Research: Solid Earth*, 126(9), e2020JB021585.
  - Hamiel, Y., Piatibratova, O., & Mizrahi, Y. (2016, March). Creep along the northern Jordan Valley section of the Dead Sea Fault. *Geophysical Research Letters*, 43(6), 2494–2501. Retrieved 2019-07-01, from https://onlinelibrary.wiley.com/doi/abs/10.1002/2016GL067913 doi: 10.1002/2016GL067913
  - Hardebeck, J. L., & Michael, A. J. (2004). Stress orientations at intermediate angles to the San Andreas Fault, California. Journal of Geophysical Research: Solid Earth, 109(B11). doi: https://doi.org/10.1029/2004JB003239
  - Hartog, R., & Schwartz, S. Y. (2001). Depth-dependent mantle anisotropy below the San Andreas fault system: Apparent splitting parameters and waveforms. Journal of Geophysical Research: Solid Earth, 106 (B3), 4155-4167. doi: https://doi.org/10.1029/2000JB900382
  - Hatzfeld, D., Karagianni, E., Kassaras, I., Kiratzi, A., Louvari, E., Lyon-Caen, H., ... Priestley, K. (2001). Shear wave anisotropy in the upper mantle beneath the Aegean related to internal deformation. *Journal of Geophysical Research: Solid Earth*, 106 (B12), 30737-30753. doi: https://doi.org/10.1029/2001JB000387
  - Hatzfeld, D., & Molnar, P. (2010). Comparisons of the kinematics and deep structures of the Zagros and Himalaya and of the Iranian and Tibetan plateaus and geodynamic implications. *Rev. Geophys.*, 48 (RG2005). doi: 10.1029/2009RG000304
  - Hayes, G. P., Moore, G. L., Portner, D. E., Hearne, M., Flamme, H., Furtney, M., & Smoczyk, G. M. (2018). Slab2, a comprehensive subduction zone geometry model. Science, 362(6410), 58–61.
  - He, P., Wen, Y., Zhong, Y., & Cai, J. (2024, 10). Nonoverlapped Sources of the Devastating 2023 Mw>6 Herat, Afghanistan, Earthquake Swarm Estimated by InSAR. Seismological Research Letters. doi: 10.1785/0220240239
  - He, X., Zhao, L.-F., Xie, X.-B., Tian, X., & Yao, Z.-X. (2021). Weak Crust in Southeast Tibetan Plateau Revealed by Lg-Wave Attenuation Tomography: Implications for Crustal Material Escape. Journal of Geophysical Research: Solid Earth, 126(3), e2020JB020748. doi: https://doi.org/10.1029/2020JB020748
  - Hearn, E. H., McClusky, S., Ergintav, S., & Reilinger, R. E. (2009). Izmit earthquake postseismic deformation and dynamics of the North Anatolian Fault Zone. *Journal of Geophysical Research: Solid Earth*, 114(B8). doi: https://doi.org/10.1029/2008JB006026
- Heidbach, O. (2005). Velocity field of the Aegean-Anatolian region from 3D finite element models. In F. Wenzel (Ed.), *Perspectives in Modern Seismology* (pp. 169–185). Berlin: Springer.
- Heidbach, O., Rajabi, M., Di Giacomo, D., Harris, J., Lammers, S., Morawietz, S.,

  Light M. O. (2025). World Stress Map 2025 [Dataset]. GFZ Data
  Services. doi: https://doi.org/10.5880/WSM.2025.002
  - Holt, W. E. (1995). Flow fields within the Tonga slab determined from the moment tensors of deep earthquakes. *Geophys. Res. Lett.*, 22, 989–992.
- Holt, W. E. (2000, 01). Correlated crust and mantle strain fields in Tibet. Geology,

28(1), 67-70. doi: 10.1130/0091-7613(2000)28(67:CCAMSF)2.0.CO;2

1423

1424

1426

1427

1428

1429

1430

1433

1434

1436

1437

1439

1440

1441

1442

1443

1445

1446

1447

1448

1449

1451

1452

1453

1454

1455

1457

1458

1459

1460

1461

1464

1465

1467

1470

1471

1472

1473

1474

1476

- Holt, W. E., Chamot-Rooke, N., Le Pichon, X., Haines, A. J., Shen-Tu, B., & Ren, J. (2000). Velocity field in Asia inferred from Quaternary fault slip rates and Global Positioning System observations. *Journal of Geophysical Research: Solid Earth*, 105(B8), 19185-19209. doi: https://doi.org/10.1029/2000JB900045
  - Holt, W. E., Ni, J. F., Wallace, T. C., & Haines, A. J. (1991). The active tectonics of the eastern Himalayan syntaxis and surrounding regions. *Journal of Geophysical Research: Solid Earth*, 96 (B9), 14595-14632. doi: https://doi.org/10.1029/91JB01021
  - Hou, Z., Liu, L., Zhang, H., Xu, B., Wang, Q., Yang, T., ... others (2024). Cenozoic eastward growth of the Tibetan Plateau controlled by tearing of the Indian slab. *Nature Geoscience*, 17(3), 255–263. doi: https://doi.org/10.1038/s41561-024-01382-9
  - Houseman, G., & England, P. (1993). Crustal thickening versus lateral expulsion in the Indian-Asian continental collision. *Journal of Geophysical Research: Solid Earth*, 98(B7), 12233-12249. doi: https://doi.org/10.1029/93JB00443
  - Huang, X., Li, Y., Shan, X., Zhao, D., Gao, Z., Gong, W., & Qu, C. (2023a, 06). InSAR Observations Reveal Variations in Shallow Creep on the Kangding Segment of the Xianshuihe Fault. Seismological Research Letters, 94(5), 2291-2300. doi: 10.1785/0220230053
  - Huang, X., Li, Y., Shan, X., Zhao, D., Gao, Z., Gong, W., & Qu, C. (2023b, 06). InSAR Observations Reveal Variations in Shallow Creep on the Kangding Segment of the Xianshuihe Fault. Seismological Research Letters, 94(5), 2291-2300. doi: 10.1785/0220230053
  - Huang, Z., Zhou, Y., & Zhang, P. (2023). Newly discovered shallow creep along the Gozha Co fault in northwestern Tibet: Spatial extent, rate and temporal evolution. *Earth and Planetary Science Letters*, 621, 118388. doi: https://doi.org/10.1016/j.epsl.2023.118388
  - Hussain, E., Hooper, A., Wright, T. J., Walters, R. J., & Bekaert, D. P. S. (2016). Interseismic strain accumulation across the central North Anatolian Fault from iteratively unwrapped InSAR measurements. *Journal of Geophysical Research:* Solid Earth, 121 (12), 9000-9019. doi: https://doi.org/10.1002/2016JB013108
  - Hussain, E., Wright, T. J., Walters, R. J., Bekaert, D. P., Lloyd, R., & Hooper, A. (2018). Constant strain accumulation rate between major earthquakes on the North Anatolian Fault. *Nature communications*, 9(1), 1392. doi: https://doi.org/10.1038/s41467-018-03739-2
  - Ismail-Zadeh, A., Adamia, S., Chabukiani, A., Chelidze, T., Cloetingh, S., Floyd, M., ... Soloviev, A. (2020). Geodynamics, seismicity, and seismic hazards of the Caucasus. *Earth-Science Reviews*, 207, 103222. doi: https://doi.org/10.1016/j.earscirev.2020.103222
  - Jackson, J., & McKenzie, D. (1984, 04). Active tectonics of the Alpine—Himalayan Belt between western Turkey and Pakistan. *Geophysical Journal International*, 77(1), 185-264. doi: 10.1111/j.1365-246X.1984.tb01931.x
  - Jenny, S., Goes, S., Giardini, D., & Kahle, H.-G. (2004). Earthquake recurrence parameters from seismic and geodetic strain rates in the eastern Mediterranean. *Geophys. J. Int.*, 157, 1331–1347.
  - Johansson, J. M., Davis, J. L., Scherneck, H.-G., Milne, G. A., Vermeer, M., Mitrovica, J. X., . . . Shapiro, I. I. (2002). Continuous GPS measurements of postglacial adjustment in Fennoscandia 1. Geodetic results. *Journal of Geophysical Research: Solid Earth*, 107(B8), ETG 3-1-ETG 3-27. doi: https://doi.org/10.1029/2001JB000400
- Johnson, K. M. (2023). BforStrain (Version 1.1.0). [Software]. Zenodo. doi: https://doi.org/10.5281/zenodo.8139779
  - Johnson, K. M. (2024). Disagreements in Geodetically Inferred Strain Rates in the

Western US With Stress Orientations and Geologic Moment Rates. *Journal of Geophysical Research: Solid Earth*, 129(4), e2023JB027472.

- Johnson, K. M., Hammond, W. C., & Weldon, I., Ray J. (2024, 01). Review of Geodetic and Geologic Deformation Models for 2023 U.S. National Seismic Hazard Model. Bulletin of the Seismological Society of America, 114(3), 1407-1436. doi: 10.1785/0120230137
- Jolivet, L., Faccenna, C., Becker, T., Tesauro, M., Sternai, P., & Bouilhol, P. (2018). Mantle Flow and Deforming Continents: From India-Asia Convergence to Pacific Subduction. *Tectonics*, 37(9), 2887-2914. doi: https://doi.org/10.1029/2018TC005036
- Jolivet, L., Faccenna, C., Huet, B., Labrousse, L., Le Pourhiet, L., Lacombe, O., ... Driussi, O. (2013). Aegean tectonics: Strain localisation, slab tearing and trench retreat. *Tectonophysics*, 597-598, 1-33. (The Aegean: a natural laboratory for tectonics Neotectonics) doi: https://doi.org/10.1016/j.tecto.2012.06.011
- Jolivet, L., Faccenna, C., & Piromallo, C. (2009). From mantle to crust: Stretching the Mediterranean. *Earth and Planetary Science Letters*, 285(1), 198-209. doi: https://doi.org/10.1016/j.epsl.2009.06.017
- Jolivet, R., Jara, J., Dalaison, M., Rouet-Leduc, B., Özdemir, A., Dogan, U., ...

  Dubernet, P. (2023). Daily to Centennial Behavior of Aseismic Slip Along the Central Section of the North Anatolian Fault. *Journal of Geophysical Research: Solid Earth*, 128(7), e2022JB026018. (e2022JB026018 2022JB026018) doi: https://doi.org/10.1029/2022JB026018
- Jónsson, S. (2008). Importance of post-seismic viscous relaxation in southern Iceland. *Nature Geoscience*, 1(2), 136–139.
- Kadirov, F., Guliyev, I., Feyzullayev, A., Safarov, R., Mammadov, S., Babayev, G., & Rashidov, T. (2014). GPS-based crustal deformations in Azerbaijan and their influence on seismicity and mud volcanism. *Izvestiya, Physics of the Solid Earth*, 50, 814–823.
- Kahle, H.-G., Straub, C., Reilinger, R., McClusky, S., King, R., Hurst, K., ... Cross,
  P. (1998). The strain rate field in the eastern Mediterranean region, estimated by repeated GPS measurements. *Tectonophys.*, 294, 237–252.
- Kaneko, Y., Fialko, Y., Sandwell, D. T., Tong, X., & Furuya, M. (2013). Interseismic deformation and creep along the central section of the North Anatolian Fault (Turkey): InSAR observations and implications for rate-and-state friction properties. *Journal of Geophysical Research: Solid Earth*, 118(1), 316-331. doi: https://doi.org/10.1029/2012JB009661
- Karakhanyan, A., Vernant, P., Doerflinger, E., Avagyan, A., Philip, H., Aslanyan, R., ... Masson, F. (2013). GPS constraints on continental deformation in the Armenian region and Lesser Caucasus. *Tectonophysics*, 592, 39-45. doi: https://doi.org/10.1016/j.tecto.2013.02.002
- Kendall, J.-M., Pilidou, S., Keir, D., Bastow, I., Stuart, G., & Ayele, A. (2006). Mantle upwellings, melt migration and the rifting of Africa: Insights from seismic anisotropy. Geol. Soc., London, Spec. Pub., 259, 55–72.
- Khorrami, F., Vernant, P., Masson, F., Nilfouroushan, F., Mousavi, Z., Nankali, H., ... others (2019). An up-to-date crustal deformation map of Iran using integrated campaign-mode and permanent GPS velocities. *Geophysical Journal International*, 217(2), 832–843.
- Kong, F., Wu, J., Liu, L., Liu, K. H., Song, J., Li, J., & Gao, S. S. (2018). Azimuthal anisotropy and mantle flow underneath the southeastern Tibetan Plateau and northern Indochina Peninsula revealed by shear wave splitting analyses. *Tectonophysics*, 747-748, 68-78. doi: https://doi.org/10.1016/j.tecto.2018.09.013
- Kostrov, V. (1974). Seismic moment and enengy of earthquakes, and the seismic flow of rock, Izv. Acad. Sci. USSR. *Phys. Solid Earth*, 1, 23–44.

Kreemer, C., Blewitt, G., & Davis, P. M. (2020, 05). Geodetic evidence for a buoyant mantle plume beneath the Eifel volcanic area, NW Europe. *Geophysical Journal International*, 222(2), 1316-1332. doi: 10.1093/gji/ggaa227

- Kreemer, C., Blewitt, G., & Hammond, W. C. (2009, 04). Geodetic constraints on contemporary deformation in the northern Walker Lane: 2. Velocity and strain rate tensor analysis. In *Late cenozoic structure and evolution of the great basin-sierra nevada transition*. Geological Society of America. doi: 10.1130/2009.2447(02)
  - Kreemer, C., Blewitt, G., & Klein, E. C. (2014). A geodetic plate motion and Global Strain Rate Model. *Geochemistry, Geophysics, Geosystems*, 15(10), 3849–3889. doi: https://doi.org/10.1002/2014GC005407
  - Kreemer, C., & Chamot-Rooke, N. (2004, 06). Contemporary kinematics of the southern Aegean and the Mediterranean Ridge. *Geophysical Journal International*, 157(3), 1377-1392. doi: 10.1111/j.1365-246X.2004.02270.x
  - Kreemer, C., Chamot-Rooke, N., & Le Pichon, X. (2004). Constraints on the evolution and vertical coherency of deformation in the Northern Aegean from a comparison of geodetic, geologic and seismologic data. Earth and Planetary Science Letters, 225(3), 329-346. doi: https://doi.org/10.1016/j.epsl.2004.06.018
  - Kreemer, C., Hammond, W. C., & Blewitt, G. (2018). A Robust Estimation of the 3-D Intraplate Deformation of the North American Plate From GPS. *Journal of Geophysical Research: Solid Earth*, 123(5), 4388-4412. doi: https://doi.org/10.1029/2017JB015257
  - Kreemer, C., Lavallée, D. A., Blewitt, G., & Holt, W. E. (2006). On the stability of a geodetic no-net-rotation frame and its implication for the International Terrestrial Reference Frame. *Geophysical Research Letters*, 33(17). doi: https://doi.org/10.1029/2006GL027058
  - Kuhlemann, J., Frisch, W., Székely, B., Dunkl, I., & Kazmer, M. (2002). Post-collisional sediment budget history of the Alps: tectonic versus climatic control. *International Journal of Earth Sciences*, 91, 818–837. doi: https://doi.org/10.1007/s00531-002-0266-y
  - Kurt, A. I., OZBAKIR, A. D., CINGOZ, A., Ergintav, S., DOGAN, U., & ÖZARPACI, S. (2023). Contemporary velocity field for Turkey inferred from combination of a dense network of long term GNSS observations. Turkish Journal of Earth Sciences, 32(3), 275–293.
  - Lazecký, M., Spaans, K., González, P. J., Maghsoudi, Y., Morishita, Y., Albino, F., ... Wright, T. J. (2020). LiCSAR: An Automatic InSAR Tool for Measuring and Monitoring Tectonic and Volcanic Activity. *Remote Sensing*, 12(15). doi: 10.3390/rs12152430
  - Le Pichon, X., & Angelier, J. (1979). The Hellenic arc and trench system: a key to the neotectonic evolution of the eastern Mediterranean area. *Tectonophysics*, 60 (1-2), 1–42.
  - Le Pichon, X., & Kreemer, C. (2010). The Miocene-to-Present Kinematic Evolution of the Eastern Mediterranean and Middle East and Its Implications for Dynamics. Annual Review of Earth and Planetary Sciences, 38 (Volume 38, 2010), 323-351. doi: https://doi.org/10.1146/annurev-earth-040809-152419
- Lev, E., Long, M. D., & van der Hilst, R. D. (2006). Seismic anisotropy in Eastern Tibet from shear wave splitting reveals changes in lithospheric deformation. Earth and Planetary Science Letters, 251(3), 293-304. doi: https://doi.org/10.1016/j.epsl.2006.09.018
- León Soto, G., Sandvol, E., Ni, J. F., Flesch, L., Hearn, T. M., Tilmann, F., ...

  Brown, L. D. (2012). Significant and vertically coherent seismic anisotropy
  beneath eastern Tibet. Journal of Geophysical Research: Solid Earth, 117(B5).
  doi: https://doi.org/10.1029/2011JB008919
  - Li, A., & Chen, C. (2006). Shear wave splitting beneath the central Tien Shan and

tectonic implications. Geophysical Research Letters, 33(22). doi: https://doi.org/10.1029/2006GL027717

- Li, X., Jónsson, S., Liu, S., Ma, Z., Castro-Perdomo, N., Cesca, S., ... Klinger, Y.
   (2024). Resolving the slip-rate inconsistency of the northern Dead Sea fault.
   Science Advances, 10(11). doi: 10.1126/sciadv.adj8408
- Li, Y., Nocquet, J.-M., Shan, X., & Jian, H. (2021). Heterogeneous Interseismic Coupling Along the Xianshuihe-Xiaojiang Fault System, Eastern Tibet.

  Journal of Geophysical Research: Solid Earth, 126(11), e2020JB021187. doi: https://doi.org/10.1029/2020JB021187
- Li, Y., Nocquet, J.-M., Shan, X., & Song, X. (2021). Geodetic Observations of Shallow Creep on the Laohushan-Haiyuan Fault, Northeastern Tibet. *Journal of Geophysical Research: Solid Earth*, 126(6), e2020JB021576. doi: https://doi.org/10.1029/2020JB021576
- Li, Y., Shan, X., Qu, C., Zhang, G., Wang, X., & Xiong, H. (2025). Slip Deficit Rate and Seismic Potential on Crustal Faults in Tibet. Geophysical Research Letters, 52(1), e2024GL112122. doi: https://doi.org/10.1029/2024GL112122
- Liang, S., Gan, W., Shen, C., Xiao, G., Liu, J., Chen, W., ... Zhou, D. (2013). Three-dimensional velocity field of present-day crustal motion of the Tibetan Plateau derived from GPS measurements. *Journal of Geophysical Research:* Solid Earth, 118(10), 5722–5732.
- Lin, Y., & Zhao, L. (2024). Upper-Mantle Anisotropy in the Southeastern Margin of Tibetan Plateau Revealed by Fullwave SKS Splitting Intensity Tomography. *Journal of Geophysical Research: Solid Earth*, 129(3), e2023JB027629. doi: https://doi.org/10.1029/2023JB027629
- Lindsey, E. O., Wang, Y., Aung, L. T., Chong, J.-H., Qiu, Q., Mallick, R., . . . others (2023). Active subduction and strain partitioning in western Myanmar revealed by a dense survey GNSS network. *Earth and Planetary Science Letters*, 622, 118384. doi: https://doi.org/10.1016/j.epsl.2023.118384
- Lippitsch, R., Kissling, E., & Ansorge, J. (2003). Upper mantle structure beneath the Alpine orogen from high-resolution teleseismic tomography. *Journal of Geophysical Research: Solid Earth*, 108(B8). doi: https://doi.org/10.1029/2002JB002016
- Liu, S., Shen, Z.-K., Bürgmann, R., & Jónsson, S. (2020, 10). Thin crème brûlée rheological structure for the Eastern California Shear Zone. Geology, 49(2), 216-221. doi: 10.1130/G47729.1
- Liu, S., Xu, X., Nocquet, J.-M., Chen, G., Tan, X., Jónsson, S., & Klinger, Y. (2025). Lower crustal thickening drives active uplift in Northern Tibet. *Earth and Planetary Science Letters*, 655, 119245. doi: https://doi.org/10.1016/j.epsl.2025.119245
- Liu, Z., Luo, H., Klinger, Y., & Wang, T. (2025). Shear Strain Evolution Spanning the 2020 Mw6.8 Elazığ and 2023 Mw7.8/Mw7.6 Kahramanmaraş Earthquake Sequence Along the East Anatolian Fault Zone. Geophysical Research Letters, 52(6), e2024GL114033. doi: https://doi.org/10.1029/2024GL114033
- Liu, Z., & Wang, T. (2023). High-Resolution Interseismic Strain Mapping From In-SAR Phase-Gradient Stacking: Application to the North Anatolian Fault With Implications for the Non-Uniform Strain Distribution Related to Coseismic Slip Distribution. Geophysical Research Letters, 50(15), e2023GL104168. doi: https://doi.org/10.1029/2023GL104168
- Loveless, J., & Meade, B. (2011). Partitioning of localized and diffuse deformation in the Tibetan Plateau from joint inversions of geologic and geodetic observations. Earth and Planetary Science Letters, 303(1), 11-24. doi: https://doi.org/10.1016/j.epsl.2010.12.014
- Luo, H., & Wang, K. (2022). Finding Simplicity in the Complexity of Postseismic Coastal Uplift and Subsidence Following Great Subduction Earthquakes. Journal of Geophysical Research: Solid Earth, 127(10), e2022JB024471. doi:

https://doi.org/10.1029/2022JB024471

- Mai, P. M., Aspiotis, T., Aquib, T. A., Cano, E. V., Castro-Cruz, D., Espindola-Carmona, A., . . . Jónsson, S. (2023, 05). The Destructive Earthquake Doublet of 6 February 2023 in South-Central Türkiye and Northwestern Syria: Initial Observations and Analyses. The Seismic Record, 3(2), 105-115. doi: 10.1785/0320230007
- Makeyeva, L., Vinnik, L., & Roecker, S. (1992). Shear-wave splitting and small-scale convection in the continental upper mantle. *Nature*, 358(6382), 144–147.
- Malinverno, A., & Ryan, W. B. F. (1986). Extension in the Tyrrhenian Sea and shortening in the Apennines as result of arc migration driven by sinking of the lithosphere. *Tectonics*, 5(2), 227-245. doi: https://doi.org/10.1029/TC005i002p00227
- Mallick, R., Hubbard, J. A., Lindsey, E. O., Bradley, K. E., Moore, J. D., Ahsan, A., ... Hill, E. M. (2020). Subduction initiation and the rise of the Shillong Plateau. Earth and Planetary Science Letters, 543, 116351. Retrieved from https://www.sciencedirect.com/science/article/pii/S0012821X20302958 doi: https://doi.org/10.1016/j.epsl.2020.116351
- Maurer, J., & Materna, K. (2023). Quantification of geodetic strain rate uncertainties and implications for seismic hazard estimates. *Geophysical Journal International*, 234(3), 2128–2142.
- McClusky, S., Balassanian, S., Barka, A., Demir, C., Ergintav, S., Georgiev, I., ... Veis, G. (2000). Global Positioning System constraints on plate kinematics and dynamics in the eastern Mediterranean and Caucasus.

  \*Journal of Geophysical Research: Solid Earth, 105 (B3), 5695-5719. doi: https://doi.org/10.1029/1999JB900351
- McClusky, S., Reilinger, R., Ogubazghi, G., Amleson, A., Healeb, B., Vernant, P., ... others (2010). Kinematics of the southern Red Sea–Afar Triple Junction and implications for plate dynamics. *Geophysical Research Letters*, 37(5).
- McKenzie, D. (1978, 10). Active tectonics of the Alpine—Himalayan belt: the Aegean Sea and surrounding regions. *Geophysical Journal International*, 55(1), 217-254. doi: 10.1111/j.1365-246X.1978.tb04759.x
- McKenzie, D., & Yılmaz, Y. (1991). Deformation and volcanism in western Turkey and the Aegean. Bulletin of the Technical University of Istanbul, 44 (1-2), 345–373.
- McKenzie, D. P. (1969). The relation between fault plane solutions for earthquakes and the directions of the principal stresses. *Bull. Seismol. Soc. Am.*, 59(2), 591–601.
- McNamara, D. E., Owens, T. J., Silver, P. G., & Wu, F. T. (1994). Shear wave anisotropy beneath the Tibetan Plateau. *Journal of Geophysical Research:* Solid Earth, 99(B7), 13655-13665. doi: https://doi.org/10.1029/93JB03406
- Meade, B. J. (2007, 01). Present-day kinematics at the India-Asia collision zone. Geology, 35(1), 81-84. doi: 10.1130/G22924A.1
- Merry, T. A. J., Bastow, I. D., Kounoudis, R., Ogden, C. S., Bell, R. E., & Jones, L. (2021). The Influence of the North Anatolian Fault and a Fragmenting Slab Architecture on Upper Mantle Seismic Anisotropy in the Eastern Mediterranean. *Geochemistry, Geophysics, Geosystems*, 22(9), e2021GC009896. doi: https://doi.org/10.1029/2021GC009896
- Mey, J., Scherler, D., Wickert, A. D., Egholm, D. L., Tesauro, M., Schildgen, T. F., & Strecker, M. R. (2016). Glacial isostatic uplift of the European Alps. *Nature Communications*, 7(1), 13382. doi: https://doi.org/10.1038/ncomms13382
- Michael, A. J. (1987). Use of focal mechanisms to determine stress: a control study. J. Geophys. Res.: Sol. Earth, 92, 357–368.
- Milne, G. A., Davis, J. L., Mitrovica, J. X., Scherneck, H.-G., Johansson, J. M., Vermeer, M., & Koivula, H. (2001). Space-Geodetic Constraints on Glacial Isostatic Adjustment in Fennoscandia. Science, 291 (5512), 2381-2385. doi:

10.1126/science.1057022

- Molnar, P., England, P., & Martinod, J. (1993). Mantle dynamics, uplift of the Tibetan Plateau, and the Indian Monsoon. *Reviews of Geophysics*, 31(4), 357-396. doi: https://doi.org/10.1029/93RG02030
- Molnar, P., & Lyon-Caen, H. (1988, 01). Some simple physical aspects of the support, structure, and evolution of mountain belts. In *Processes in Continental Lithospheric Deformation*. Geological Society of America. doi: 10.1130/SPE218-p179
- Molnar, P., & Tapponnier, P. (1975). Cenozoic Tectonics of Asia: Effects of a Continental Collision. *Science*, 189(4201), 419-426. doi: 10.1126/science.189.4201.419
- Mumladze, T., Forte, A. M., Cowgill, E. S., Trexler, C. C., Niemi, N. A., Burak Yıkılmaz, M., & Kellogg, L. H. (2015). Subducted, detached, and torn slabs beneath the Greater Caucasus. *GeoResJ*, 5, 36-46. doi: https://doi.org/10.1016/j.grj.2014.09.004
- Nicolas, A., & Christensen, N. I. (1987). Formation of Anisotropy in Upper Mantle Peridotites A Review. In *Composition, Structure and Dynamics of the Lithosphere-Asthenosphere System* (p. 111-123). American Geophysical Union (AGU). doi: https://doi.org/10.1029/GD016p0111
- Nocquet, J.-M. (2012). Present-day kinematics of the Mediterranean: A comprehensive overview of GPS results. *Tectonophysics*, 579, 220–242.
- Nocquet, J.-M., Sue, C., Walpersdorf, A., Tran, T., Lenôtre, N., Vernant, P., ... others (2016). Present-day uplift of the western Alps. *Scientific reports*, 6(1), 28404. doi: https://doi.org/10.1038/srep28404
- Nucci, R., Serpelloni, E., Faenza, L., Garcia, A., & Belardinelli, M. E. (2025). Geodetic Strain Rates and Seismicity Rates Along the Apennines (Italy). Journal of Geophysical Research: Solid Earth, 130(2), e2024JB029848. doi: https://doi.org/10.1029/2024JB029848
- Okazaki, T., Fukahata, Y., & Nishimura, T. (2021). Consistent estimation of strainrate fields from GNSS velocity data using basis function expansion with ABIC. Earth, Planets and Space, 73, 1–22.
- Ou, Q., Daout, S., Weiss, J. R., Shen, L., Lazecký, M., Wright, T. J., & Parsons, B. E. (2022). Large-Scale Interseismic Strain Mapping of the NE Tibetan Plateau From Sentinel-1 Interferometry. Journal of Geophysical Research: Solid Earth, 127(6), e2022JB024176. doi: https://doi.org/10.1029/2022JB024176
- Özarpacı, S., Doğan, U., Ergintav, S., Çakır, Z., Özdemir, A., Floyd, M., & Reilinger, R. (2021). Present GPS velocity field along 1999 Izmit rupture zone: evidence for continuing afterslip 20 yr after the earthquake. Geophysical Journal International, 224(3), 2016–2027.
- Özbey, V., Sengör, A. M. C., Henry, P., Özeren, M. S., Haines, A. J., Klein, E. C., ... Öğretmen, N. (2024, 8). Kinematics of the Kahramanmaraş triple junction and of Cyprus: evidence of shear partitioning. *BSGF Earth Sciences Bulletin*, 195, 15. doi: 10.1051/bsgf/2024012
- Ozdemir, S., & Karshoğlu, M. O. (2019). Soft clustering of GPS velocities from a homogeneous permanent network in Turkey. *Journal of Geodesy*, 93(8), 1171–1195.
- Özkan, A., Yavaşoğlu, H. H., & Masson, F. (2023). Present-day strain accumulations and fault kinematics at the Hatay Triple Junction using new geodetic constraints. *Tectonophysics*, 854, 229819.
- Pagani, C., Bodin, T., Métois, M., & Lasserre, C. (2021). Bayesian estimation of surface strain rates from global navigation satellite system measurements: Application to the southwestern United States. *Journal of Geophysical Research:* Solid Earth, 126(6), e2021JB021905.
- Palano, M. (2014, 12). On the present-day crustal stress, strain-rate fields and man-

tle anisotropy pattern of Italy. Geophysical Journal International, 200(2), 969-985. doi: 10.1093/gji/ggu451

1756

1757

1758

1759

1760

1763

1765

1766

1769

1770

1771

1772

1773

1775

1776

1778

1781

1782

1783

1787

1788

1789

1790

1792

1793

1794

1795

1796

1799

1800

1801

1802

1803

1805

1806

- Papaleo, E., Cornwell, D., & Rawlinson, N. (2018). Constraints on North Anatolian Fault Zone Width in the Crust and Upper Mantle From S Wave Teleseismic Tomography. *Journal of Geophysical Research: Solid Earth*, 123(4), 2908-2922. doi: https://doi.org/10.1002/2017JB015386
- Paul, A., Karabulut, H., Mutlu, A. K., & Salaün, G. (2014). A comprehensive and densely sampled map of shear-wave azimuthal anisotropy in the Aegean-Anatolia region. Earth and Planetary Science Letters, 389, 14-22. doi: https://doi.org/10.1016/j.epsl.2013.12.019
- Peltzer, G., & Tapponnier, P. (1988). Formation and evolution of strike-slip faults, rifts, and basins during the India-Asia Collision: An experimental approach. *Journal of Geophysical Research: Solid Earth*, 93 (B12), 15085-15117. doi: https://doi.org/10.1029/JB093iB12p15085
- Perry, M., Kakar, N., Ischuk, A., Metzger, S., Bendick, R., Molnar, P., & Mohadjer, S. (2019). Little geodetic evidence for localized Indian subduction in the Pamir-Hindu Kush of Central Asia. *Geophysical Research Letters*, 46(1), 109–118.
- Piña-Valdés, J., Socquet, A., Beauval, C., Doin, M.-P., D'Agostino, N., & Shen, Z.-K. (2022). 3D GNSS velocity field sheds light on the deformation mechanisms in Europe: Effects of the vertical crustal motion on the distribution of seismicity. *Journal of Geophysical Research: Solid Earth*, 127(6), e2021JB023451.
- Podolefsky, N. S., Zhong, S., & McNamara, A. K. (2004). The anisotropic and rheological structure of the oceanic upper mantle from a simple model of plate shear. *Geophys. J. Int.*, 158, 287–296.
- Qiao, X., & Zhou, Y. (2021). Geodetic imaging of shallow creep along the Xianshuihe fault and its frictional properties. *Earth and Planetary Science Letters*, 567, 117001. doi: https://doi.org/10.1016/j.epsl.2021.117001
- Read, H. H., & Watson, J. (1975). Introduction to Geology: Earth History. Part II: Later Stages of Earth History. Springer. doi: 10.1007/978-1-349-15613-9
- Rebaï, S., Philip, H., Dorbath, L., Borissoff, B., Haessler, H., & Cisternas, A. (1993). Active tectonics in the Lesser Caucasus: Coexistence of compressive and extensional structures. *Tectonics*, 12(5), 1089-1114. doi: https://doi.org/10.1029/93TC00514
- Reguzzoni, M., & Sampietro, D. (2015). GEMMA: An Earth crustal model based on GOCE satellite data. International Journal of Applied Earth Observation and Geoinformation, 35, 31-43. (GOCE earth science applications and models(Based on the ESA GOCE solid earth workshop, 16-17 October 2012)) doi: https://doi.org/10.1016/j.jag.2014.04.002
- Reilinger, R., & McClusky, S. (2011, 09). Nubia–Arabia–Eurasia plate motions and the dynamics of Mediterranean and Middle East tectonics. *Geophysical Journal International*, 186(3), 971-979. doi: 10.1111/j.1365-246X.2011.05133.x
- Reilinger, R., McClusky, S., Vernant, P., Lawrence, S., Ergintav, S., Cakmak, R., ... Karam, G. (2006). GPS constraints on continental deformation in the Africa-Arabia-Eurasia continental collision zone and implications for the dynamics of plate interactions. *Journal of Geophysical Research: Solid Earth*, 111 (B5). Retrieved from https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005JB004051 doi: 10.1029/2005JB004051
- Replumaz, A., & Tapponnier, P. (2003). Reconstruction of the deformed collision zone Between India and Asia by backward motion of lithospheric blocks. *Journal of Geophysical Research: Solid Earth*, 108(B6). doi: https://doi.org/10.1029/2001JB000661
- Ritter, J. R., Jordan, M., Christensen, U. R., & Achauer, U. (2001). A mantle plume below the Eifel volcanic fields, Germany. Earth Planet. Sci. Lett., 186, 7–14.

Rodriguez, M., Sakellariou, D., Gorini, C., Janin, A., d'Acremont, E., Pourhiet,

L. L., ... others (2023). Evolution of the North Anatolian Fault from a
diffuse to a localized shear zone in the North Aegean Sea during the PlioPleistocene. Geophysical Journal International, 235(3), 2614–2639. doi:
https://doi.org/10.1093/gji/ggad364

- Royden, L. H. (1993). The tectonic expression slab pull at continental convergent boundaries. *Tectonics*, 12(2), 303-325. doi: https://doi.org/10.1029/92TC02248
  - Royden, L. H., Burchfiel, B. C., King, R. W., Wang, E., Chen, Z., Shen, F., & Liu, Y. (1997). Surface Deformation and Lower Crustal Flow in Eastern Tibet. *Science*, 276(5313), 788-790. doi: 10.1126/science.276.5313.788
  - Saleh, M., & Becker, M. (2014). A new velocity field from the analysis of the Egyptian Permanent GPS Network (EPGN). Arabian Journal of Geosciences, 7, 4665–4682.
  - Sandwell, D. T. (2022). Elastic Solutions for Strike-Slip Faulting. In *Advanced Geodynamics: The Fourier Transform Method* (p. 145–168). Cambridge University Press.
  - Sandwell, D. T., & Wessel, P. (2016). Interpolation of 2-D vector data using constraints from elasticity. *Geophysical Research Letters*, 43(20), 10–703.
  - Savage, J., Gan, W., & Svarc, J. (2001). Strain accumulation and rotation in the Eastern California Shear Zone. Journal of Geophysical Research: Solid Earth, 106 (B10), 21995–22007.
  - Savage, J., & Prescott, W. (1978). Asthenosphere readjustment and the earthquake cycle. *Journal of Geophysical Research: Solid Earth*, 83(B7), 3369–3376.
  - Savage, J. C. (1983). Strain accumulation in western United States. Annual review of earth and planetary sciences, 11(1), 11–41.
  - Savage, J. C., & Burford, R. O. (1973, February). Geodetic determination of relative plate motion in central California. Journal of Geophysical Research, 78(5), 832–845. Retrieved 2018-07-17, from http://doi.wiley.com/10.1029/JB078i005p00832 doi: 10.1029/JB078i005p00832
  - Savage, M. K. (1999). Seismic anisotropy and mantle deformation: What have we learned from shear wave splitting? *Reviews of Geophysics*, 37(1), 65-106. doi: https://doi.org/10.1029/98RG02075
  - Savage, M. K., Fischer, K. M., & Hall, C. E. (2004, 01). Strain modelling, seismic anisotropy and coupling at strike-slip boundaries: applications in New Zealand and the San Andreas fault. In *Vertical Coupling and Decoupling in the Lithosphere*. Geological Society of London. doi: 10.1144/GSL.SP.2004.227.01.02
  - Schulte-Pelkum, V., Becker, T. W., Behr, W. M., & Miller, M. S. (2021). Tectonic inheritance during plate boundary evolution in southern California constrained from seismic anisotropy. *Geochem.*, *Geophys.*, *Geosys.*, 22, e2021GC010099. doi: 10.1029/2021GC010099
  - Searle, M., Elliott, J., Phillips, R., & Chung, S.-L. (2011). Crustal—lithospheric structure and continental extrusion of Tibet. *Journal of the Geological Society*, 168(3), 633-672. doi: 10.1144/0016-76492010-139
- Serpelloni, E., Cavaliere, A., Martelli, L., Pintori, F., Anderlini, L., Borghi, A., . . . others (2022). Surface velocities and strain-rates in the Euro-Mediterranean region from massive GPS data processing. Frontiers in Earth Science, 10, 907897.
- Serpelloni, E., Faccenna, C., Spada, G., Dong, D., & Williams, S. D. P. (2013). Vertical GPS ground motion rates in the Euro-Mediterranean region: New evidence of velocity gradients at different spatial scales along the Nubia-Eurasia plate boundary. *Journal of Geophysical Research: Solid Earth*, 118(11), 6003-6024. doi: https://doi.org/10.1002/2013JB010102
- Shaw, M., & Pysklywec, R. (2007). Anomalous uplift of the Apennines and subsidence of the Adriatic: The result of active mantle flow? *Geophys. Res. Lett.*,

34 (L04311). doi: 10.1029/2006GL028337

- Shen, F., Royden, L. H., & Burchfiel, B. C. (2001). Large-scale crustal deformation of the Tibetan Plateau. *Journal of Geophysical Research: Solid Earth*, 106 (B4), 6793-6816. doi: https://doi.org/10.1029/2000JB900389
- Shen, Z., & Bird, P. (2022, 09). NeoKinema Deformation Model for the 2023 Update to the U.S. National Seismic Hazard Model. Seismological Research Letters, 93(6), 3037-3052. doi: 10.1785/0220220179
- Shen, Z.-K., Jackson, D. D., & Ge, B. X. (1996). Crustal deformation across and beyond the Los Angeles basin from geodetic measurements. *Journal of Geophysical Research: Solid Earth*, 101(B12), 27957–27980.
- Shen, Z.-K., Lü, J., Wang, M., & Bürgmann, R. (2005). Contemporary crustal deformation around the southeast borderland of the Tibetan Plateau. *Journal of Geophysical Research: Solid Earth*, 110 (B11). doi: https://doi.org/10.1029/2004JB003421
- Shen, Z.-K., Wang, M., Zeng, Y., & Wang, F. (2015). Optimal interpolation of spatially discretized geodetic data. *Bulletin of the Seismological Society of America*, 105(4), 2117–2127.
- Silver, P. G. (1996). SEISMIC ANISOTROPY BENEATH THE CONTINENTS: Probing the Depths of Geology [Journal Article]. Annual Review of Earth and Planetary Sciences, 24 (Volume 24, 1996), 385-432. doi: https://doi.org/10.1146/annurev.earth.24.1.385
- Silver, P. G., & Chan, W. W. (1991). Shear wave splitting and subcontinental mantle deformation. Journal of Geophysical Research: Solid Earth, 96 (B10), 16429-16454. doi: https://doi.org/10.1029/91JB00899
- Silver, P. G., & Holt, W. E. (2002). The mantle flow field beneath Western North America. *Science*, 295, 1054–1057.
- Sokhadze, G., Floyd, M., Godoladze, T., King, R., Cowgill, E., Javakhishvili, Z., . . . Reilinger, R. (2018). Active convergence between the Lesser and Greater Caucasus in Georgia: Constraints on the tectonic evolution of the Lesser–Greater Caucasus continental collision.

  Earth and Planetary Science Letters, 481, 154–161.
- Sol, S., Meltzer, A., Burgmann, R., Van der Hilst, R., King, R., Chen, Z., ... others (2007). Geodynamics of the southeastern Tibetan Plateau from seismic anisotropy and geodesy. *Geology*, 35(6), 563–566. doi: 10.1130/G23408A.1
- Stamps, D. S., Saria, E., & Kreemer, C. (2018). A geodetic strain rate model for the East African Rift System. *Scientific reports*, 8(1), 732.
- Steffen, R., Steffen, H., Kenyeres, A., Nilsson, T., & Lidberg, M. (2025, 02). Eu-VeM2022—a European three-dimensional GNSS velocity model based on least-squares collocation. *Geophysical Journal International*, 241(1), 437-453. doi: 10.1093/gji/ggaf052
- Sternai, P., Sue, C., Husson, L., Serpelloni, E., Becker, T. W., Willett, S. D., ... Castelltort, S. (2019). Present-day uplift of the European Alps: Evaluating mechanisms and models of their relative contributions. *Earth-Science Reviews*, 190, 589-604. doi: https://doi.org/10.1016/j.earscirev.2019.01.005
- Sue, C., Thouvenot, F., Fréchet, J., & Tricart, P. (1999). Widespread extension in the core of the western Alps revealed by earthquake analysis. *Journal of Geo*physical Research: Solid Earth, 104 (B11), 25611-25622. doi: https://doi.org/10 .1029/1999JB900249
- Sun, H., Chang, L., Song, X., & Wang, X. (2025). Crustal deformation across the southeastern flank of the eastern Himalayan syntaxis from 3D velocity and anisotropic structures. Earth and Planetary Science Letters, 119230. doi: https://doi.org/10.1016/j.epsl.2025.119230
- Tape, C., Musé, P., Simons, M., Dong, D., & Webb, F. (2009). Multiscale estimation of GPS velocity fields. *Geophysical Journal International*, 179(2), 945–971.
- Tapponnier, P., Peltzer, G., Le Dain, A. Y., Armijo, R., & Cobbold, P. (1982, 12).

Propagating extrusion tectonics in Asia: New insights from simple experiments with plasticine. Geology, 10(12), 611-616. doi: 10.1130/0091-7613(1982)10/611: PETIAN $\rangle$ 2.0.CO;2

- Tapponnier, P., Zhiqin, X., Roger, F., Meyer, B., Arnaud, N., Wittlinger, G., & Jingsui, Y. (2001). Oblique Stepwise Rise and Growth of the Tibet Plateau. *Science*, 294(5547), 1671-1677. doi: 10.1126/science.105978
- Tatar, O., Poyraz, F., Gürsoy, H., Cakir, Z., Ergintav, S., Akpınar, Z., . . . others (2012). Crustal deformation and kinematics of the Eastern Part of the North Anatolian Fault Zone (Turkey) from GPS measurements. *Tectonophysics*, 518, 55–62.
- Taylor, G., Rost, S., & Houseman, G. (2016). Crustal imaging across the North Anatolian Fault Zone from the autocorrelation of ambient seismic noise. *Geophysical Research Letters*, 43(6), 2502-2509. doi: https://doi.org/10.1002/2016GL067715
- Ten Brink, U., Schoenberg, N., Kovach, R., & Ben-Avraham, Z. (1990). Uplift and a possible moho offset across the Dead Sea transform. *Tectonophysics*, 180(1), 71-85. doi: https://doi.org/10.1016/0040-1951(90)90373-G
- Thatcher, W. (2007). Microplate model for the present-day deformation of Tibet. Journal of Geophysical Research: Solid Earth, 112(B1). doi: https://doi.org/10.1029/2005JB004244
- Thatcher, W. (2009). How the Continents Deform: The Evidence From Tectonic Geodesy\* [Journal Article]. Annual Review of Earth and Planetary Sciences, 37(Volume 37, 2009), 237-262. doi: https://doi.org/10.1146/annurev.earth.031208.100035
- Tian, D., Uieda, L., Leong, W. J., Fröhlich, Y., Schlitzer, W., Grund, M., ...
  Wessel, P. (2024). PyGMT: A Python interface for the Generic Mapping
  Tools (Version 0.12.0) [Software]. Zenodo. doi: https://doi.org/10.5281/
  zenodo.11062720
- Timsina, P., Hearn, T. M., & Ni, J. F. (2024). Crust and Mantle Flow From Central Tibetan Plateau to the Indo-Burma Subduction Zone. *Journal of Geophysical Research: Solid Earth*, 129(10), e2023JB027540. doi: https://doi.org/10.1029/2023JB027540
- Tong, X., & Chen, S. (2024, 08). Temporal and Spatial Variation of Fault Creep along the Xianshuihe Fault from InSAR Stacking. Geophysical Journal International, ggae310. doi: 10.1093/gji/ggae310
- Tosi, L., Teatini, P., & Strozzi, T. (2013). Natural versus anthropogenic subsidence of Venice. Scientific reports, 3(1), 2710. doi: https://doi.org/10.1038/srep02710
- Townend, J., & Zoback, M. D. (2004). Regional tectonic stress near the San Andreas fault in central and southern California. *Geophysical Research Letters*, 31(15). doi: https://doi.org/10.1029/2003GL018918
- Townend, J., & Zoback, M. D. (2006). Stress, strain, and mountain building in central Japan. *Journal of Geophysical Research: Solid Earth*, 111(B3). doi: https://doi.org/10.1029/2005JB003759
- van Gelder, I., Willingshofer, E., Sokoutis, D., & Cloetingh, S. (2017). The interplay between subduction and lateral extrusion: A case study for the European Eastern Alps based on analogue models. *Earth and Planetary Science Letters*, 472, 82-94. doi: https://doi.org/10.1016/j.epsl.2017.05.012
- Vardić, K., Clarke, P. J., & Whitehouse, P. L. (2022). A GNSS velocity field for crustal deformation studies: The influence of glacial isostatic adjustment on plate motion models. *Geophysical Journal International*, 231(1), 426–458.
- Vauchez, A., Tommasi, A., & Mainprice, D. (2012). Faults (shear zones) in the Earth's mantle. *Tectonophys.*, 558, 1–27.
- Vilotte, J. P., Madariaga, R., Daigniéres, M., & Zienkiewicz, O. (1986, 02). Numerical study of continental collision: influence of buoyancy forces and an

initial stiff inclusion. Geophysical Journal International, 84(2), 279-310. doi: 10.1111/j.1365-246X.1986.tb04357.x

1976

1977

1978

1979

1980

1983

1984

1985

1986

1989

1990

1991

1992

1995

1996

1997

1998

1999

2001

2002

2004

2007

2008

2009

2010

2011

2013

2014

2015

2016

2017

2019

2020

2021

2022

2023

2025

- Viltres, R., Doubre, C., Doin, M.-P., & Masson, F. (2025, 02). Assismic creep and strain partitioning accommodating the Nubia-Eurasia oblique convergence in northern Africa from InSAR analysis. *Geology*. doi: 10.1130/G53117.1
- Viltres, R., Jónsson, S., Alothman, A. O., Liu, S., Leroy, S., Masson, F., ... Reilinger, R. (2022). Present-Day Motion of the Arabian Plate. *Tectonics*, 41(3), e2021TC007013.
- Viltres, R., Jónsson, S., Ruch, J., Doubre, C., Reilinger, R., Floyd, M., & Ogubazghi, G. (2020). Kinematics and deformation of the southern Red Sea region from GPS observations. *Geophysical Journal International*, 221(3), 2143–2154.
- Vinnik, L. P., Makeyeva, L. I., Milev, A., & Usenko, A. Y. (1992, 12). Global patterns of azimuthal anisotropy and deformations in the continental mantle. *Geophysical Journal International*, 111(3), 433-447. doi: 10.1111/j.1365-246X.1992.tb02102.x
- Walker, K., Bokelmann, G., Klemperer, S., & Bock, G. (2005). Shear-wave splitting around the Eifel hotspot: evidence for a mantle upwelling. *Geophys. J. Int.*, 163, 962–980.
- Wang, C.-Y., Flesch, L. M., Silver, P. G., Chang, L.-J., & Chan, W. W. (2008, 05).
  Evidence for mechanically coupled lithosphere in central Asia and resulting implications. *Geology*, 36(5), 363-366. doi: 10.1130/G24450A.1
- Wang, L., & Barbot, S. (2023). Three-dimensional kinematics of the India–Eurasia collision. Communications Earth & Environment, 4(1), 164.
- Wang, L., Wang, R., Roth, F., Enescu, B., Hainzl, S., & Ergintav, S. (2009, 09). Afterslip and viscoelastic relaxation following the 1999 M 7.4 İzmit earth-quake from GPS measurements. Geophysical Journal International, 178(3), 1220-1237. doi: 10.1111/j.1365-246X.2009.04228.x
- Wang, M., & Shen, Z.-K. (2020). Present-day crustal deformation of continental China derived from GPS and its tectonic implications. *Journal of Geophysical Research: Solid Earth*, 125(2), e2019JB018774.
- Watson, A. R., Elliott, J. R., Lazecký, M., Maghsoudi, Y., McGrath, J. D., & Walters, R. J. (2024). An InSAR-GNSS Velocity Field for Iran. *Geophysical Research Letters*, 51(10), e2024GL108440. doi: https://doi.org/10.1029/2024GL108440
- Wedmore, L. N., Biggs, J., Floyd, M., Fagereng, Å., Mdala, H., Chindandali, P., ... Mphepo, F. (2021). Geodetic constraints on cratonic microplates and broad strain during rifting of thick Southern African lithosphere. Geophysical Research Letters, 48(17), e2021GL093785.
- Weertman, J., & Weertman, J. (1964). Elementary dislocation theory. Macmillan. Retrieved from https://books.google.com.sa/books?id=Ok1RAAAAMAAJ
- Weiss, J. R., Walters, R. J., Morishita, Y., Wright, T. J., Lazecky, M., Wang, H., ... others (2020). High-resolution surface velocities and strain for Anatolia from Sentinel-1 InSAR and GNSS data. *Geophysical Research Letters*, 47(17). doi: https://doi.org/10.1029/2020GL087376
- Wessel, P., Luis, J. F., Uieda, L., Scharroo, R., Wobbe, F., Smith, W. H. F., & Tian, D. (2019). The Generic Mapping Tools Version 6. *Geochemistry, Geophysics, Geosystems*, 20(11), 5556-5564. doi: https://doi.org/10.1029/2019GC008515
- Wolfe, C. J., & Solomon, S. C. (1998). Shear-Wave Splitting and Implications for Mantle Flow Beneath the MELT Region of the East Pacific Rise. Science, 280 (5367), 1230-1232. doi: 10.1126/science.280.5367.1230
- Wortel, M. J. R., & Spakman, W. (2000). Subduction and Slab Detachment in the Mediterranean-Carpathian Region. Science, 290(5498), 1910-1917. doi: 10.1126/science.290.5498.1910
- <sup>2027</sup> Wu, D.-L., Ge, W.-P., Liu, S.-Z., Yuan, D.-Y., Zhang, B., & Wei, C.-M. (2024).

Present-day 3D crustal deformation of the Northeastern Tibetan Plateau from space geodesy. Geophysical Research Letters, 51(4), e2023GL106143. doi: https://doi.org/10.1029/2023GL106143

2031

2032

2033

2035

2038

2039

2040

2041

2042

2044

2045

2046

2047

2050

2051

2052

2053

2056

2057

2058

2059

2060

2062

2063

2064

2065

2068

2069

2070

2071

2072

2074

2075

2077

2078

2080

2081

- Wu, Y., Zheng, Z., Nie, J., Chang, L., Su, G., Yin, H., ... others (2022). High-precision vertical movement and three-dimensional deformation pattern of the Tibetan plateau. *Journal of Geophysical Research: Solid Earth*, 127(4), e2021JB023202. doi: https://doi.org/10.1029/2021JB023202
- Yamamoto, R., Kido, M., Ohta, Y., Takahashi, N., Yamamoto, Y., Pinar, A., ... Kaneda, Y. (2019). Seafloor Geodesy Revealed Partial Creep of the North Anatolian Fault Submerged in the Sea of Marmara. Geophysical Research Letters, 46(3), 1268-1275. doi: https://doi.org/10.1029/2018GL080984
- Yang, W., & Hauksson, E. (2013, 04). The tectonic crustal stress field and style of faulting along the Pacific North America Plate boundary in Southern California. *Geophysical Journal International*, 194(1), 100-117. doi: 10.1093/gji/ggt113
- Yeck, W. L., Hatem, A. E., Goldberg, D. E., Barnhart, W. D., Jobe, J. A. T., Shelly, D. R., . . . Earle, P. S. (2023, 12). Rapid Source Characterization of the 2023 Mw 6.8 Al Haouz, Morocco, Earthquake. The Seismic Record, 3(4), 357-366. doi: 10.1785/0320230040
- Yin, A., & Harrison, T. M. (2000). Geologic Evolution of the Himalayan-Tibetan Orogen. Annual Review of Earth and Planetary Sciences, 28 (Volume 28, 2000), 211-280. doi: https://doi.org/10.1146/annurev.earth.28.1.211
- Zelenin, E., Bachmanov, D., Garipova, S., Trifonov, V., & Kozhurin, A. (2021). The database of the active faults of Eurasia (AFEAD): Ontology and design behind the continental-scale dataset. *Earth System Science Data Discussions*, 2021, 1–20. doi: https://doi.org/10.5194/essd-14-4489-2022
- Zeng, Y. (2022). GPS velocity field of the western United States for the 2023 National Seismic Hazard Model Update. Seismological Society of America, 93(6), 3121–3134. doi: https://doi.org/10.1785/0220220180
- Zhang, J., ze Wen, X., ling Cao, J., Yan, W., lin Yang, Y., & Su, Q. (2018). Surface creep and slip-behavior segmentation along the northwestern Xianshuihe fault zone of southwestern China determined from decades of fault-crossing short-baseline and short-level surveys. *Tectonophysics*, 722, 356-372. doi: https://doi.org/10.1016/j.tecto.2017.11.002
- Zhang, S., & Karato, S.-i. (1995). Lattice preferred orientation of olivine aggregates deformed in simple shear. *Nature*, 375 (6534), 774–777. doi: https://doi.org/10.1038/375774a0
- Zhang, Y. G., Zheng, W. J., Wang, Y. J., Zhang, D. L., Tian, Y. T., Wang, M., ... Zhang, P. Z. (2018). Contemporary Deformation of the North China Plain From Global Positioning System Data. Geophysical Research Letters, 45(4), 1851-1859. doi: https://doi.org/10.1002/2017GL076599
- Zheng, G., Wang, H., Wright, T. J., Lou, Y., Zhang, R., Zhang, W., ... Wei, N. (2017). Crustal deformation in the India-Eurasia collision zone from 25 years of GPS measurements. *Journal of Geophysical Research: Solid Earth*, 122(11), 9290–9312. doi: https://doi.org/10.1002/2017JB014465
- Zubovich, A., Schöne, T., Metzger, S., Mosienko, O., Mukhamediev, S., Sharshebaev, A., & Zech, C. (2016). Tectonic interaction between the Pamir and Tien Shan observed by GPS. *Tectonics*, 35(2), 283–292. doi: https://doi.org/10.1002/2015TC004055
- Çakir, Z., Ergintav, S., Özener, H., Dogan, U., Akoglu, A. M., Meghraoui, M., & Reilinger, R. (2012, 12). Onset of aseismic creep on major strike-slip faults. Geology, 40(12), 1115-1118. doi: 10.1130/G33522.1
- Özalaybey, S., & Savage, M. K. (1995). Shear-wave splitting beneath western United States in relation to plate tectonics. *Journal of Geophysical Research: Solid Earth*, 100(B9), 18135-18149. doi: https://doi.org/10.1029/95JB00715

Sengör, A. (1986). The dual nature of the Alpine-Himalayan system: Progress, problems and prospects. Tectonophysics, 127(3), 177-195. (Tectonics of the Eurasian Fold Belts) doi: https://doi.org/10.1016/0040-1951(86)90060-0
Sengör, A., Görür, N., & Şaroğlu, F. (1985, 01). Strike-Slip Faulting and Related Basin Formation in Zones of Tectonic Escape: Turkey as a Case Study. In Strike-Slip Deformation, Basin Formation, and Sedimentation. SEPM Society for Sedimentary Geology. doi: 10.2110/pec.85.37.0211

Figure :
----------




Figure 2
----------






Figure 3.
-----------



IIguie 4	Fi	gu	re	4.
----------	----	----	----	----



Figu	re 5.
------	-------





Figure 6.		



Fi	g	u	r	e	7	




Figure 8.				




Figure 9
----------

