

Consolidation and Fabric in Ursa Mudstones

Ruarri Day-Stirrat: Bureau of Economic Geology

ABSTRACT

We use core material to illuminate the micro-scale fabric of clay-rich mudstones during compaction in shallow buried sediments cored as part of Integrated Ocean Drilling Program Expedition 308, Ursa Basin, Gulf of Mexico. This study is the first step in developing a comprehensive model for understanding the anisotropic behavior of mudstones. Mudstones of consistent composition, grain size and gamma-ray log response decrease in porosity from 80% at the mudline to 37% at 600m of vertical burial at Ursa. We use Argon-ion milled surfaces to image this pore evolution over a vertical effective stress range of 0.24 to 4.05 MPa. We find an elongation of the pore shape and quantify the preferential loss of large pores, primarily associated with larger grains, and decrease in mean pore size over this stress range. This soil fabric change is accompanied by a small clay mineral preferred orientation increase, also a function of effective stress, which we document in the shallow section for the first time.

CLICK ON IMAGE FOR LARGER VIEW




Fig. 1: Illite-smectite preferred orientation of Ursa mudstones as a function of vertical effective stress. Straight line is a simple linear model predicting Illite-smectite preferred orientation from vertical effective stress. Curved line incorporates void ratio into the prediction of preferred orientation.




Fig. 2: Close off model for pore throat size distribution




Fig. 1: Illite-smectite preferred orientation of Ursa mudstones as a function of vertical effective stress. Straight line is a simple linear model predicting Illite-smectite preferred orientation from vertical effective stress. Curved line incorporates void ratio into the prediction of preferred orientation.

[Back](#)




Fig. 2: Close off model for pore throat size distribution